

Smoking may impact survival after a breast cancer diagnosis

June 24 2015



Credit: Vera Kratochvil/public domain

Researchers have found that smoking may increase the risk of dying early in premenopausal women with breast cancer.

In a prospective study of 848 women with [breast cancer](#) who were followed for a median of 6.7 years, premenopausal women who smoked for more than 21.5 years had a 3.1-times higher risk of dying from any

cause as well as a 3.4-times higher risk of dying from breast cancer. These links were not apparent among post-menopausal women.

There was also some suggestion that the increased risks seen in [premenopausal women](#) were especially relevant to women whose cancers expressed both the estrogen receptor and the progesterone receptor.

"Overall, this work is monumental in advising patients about how their smoking might affect their outcome," said Dr. Yuko Minami, co-author of the Cancer Science study. "Hopefully this paper will serve to reduce the number of [breast cancer patients](#) who continue to smoke."

More information: "Smoking and survival after breast cancer diagnosis in Japanese women: A prospective cohort study." [DOI: 10.1111/cas.12716](#)

Provided by Wiley

Citation: Smoking may impact survival after a breast cancer diagnosis (2015, June 24) retrieved 25 April 2024 from <https://medicalxpress.com/news/2015-06-impact-survival-breast-cancer-diagnosis.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--