

## The majority of amateur athletes undergoing hypoxic training are not advised by specialists

June 26 2015



An athlete doing hypoxic training with professional monitoring. Credit: Jesús Álvarez-Herms



Physical performance after periods of hypoxic training in low-oxygen conditions has become a matter of growing controversy within the scientific community. An international study, with the help of Spanish researchers, compared professional and amateur athletes' knowledge and understanding of this type of training. According to the results, just 25% of amateurs are assessed and monitored by specialists.

The most popular method of training in low-oxygen conditions is known as intermittent hypoxic training. This consists of creating natural or artificial conditions that result in increased oxygen deficiency in the subject either when they are exercising or at rest.

The technique has proved to be efficient in improving oxygen transfer in some athletes, thereby improving their physical performances. However, this cannot be extrapolated to all the cases in which this type of training has been used.

Therefore, a study published in the journal *Physiology & Behaviour* sought to compare professional and amateur athletes' use, methodology, <u>knowledge</u> and understanding of hypoxic training.

"Medical and scientific monitoring and regulation of the physiological responses to exposure to this type of training is not all that widespread among amateurs. Only 25% make use of it as opposed to 98% of professionals," lead researcher Jesús Álvarez-Herms, of the UB, explained to SINC.

According to the authors, the data corresponds to the fact that professional athletes bear certain aspects very much in mind, such as nutrition to control breathlessness and avoid possible increased deficiencies. For instance, when professional athletes undergo hypoxic training, they seriously consider using iron supplements. In contrast, amateurs, for the most part, follow programmes on their own, without



specialist monitoring.

As a <u>matter</u> of fact, the lack of monitoring of their hypoxic training can lead to certain health problems for the athletes. "The risk is connected with individuals' low tolerance to altitude," says Álvarez-Herms, which relates to an increase in breathlessness, the onset of anaemia and loss of muscle mass as the possible adverse effects of hypoxia.

The differences in regulation between professionals and amateurs suggest that the latter group follow hypoxic training programmes "in which they are responsible for themselves or that are assessed by people who are not fully trained in the area. The consequences of this are increased health risks and the possibility that their efforts will prove to be ineffective," added the scientist.

## **Professional athletes are more cautious with their results**

In order to conduct the study, the experts drew up a questionnaire with 17 questions, which was completed by a total of 203 athletes—95 professionals and 108 amateurs—from different sporting disciplines (cycling, triathlon and endurance running) during the 2013-2014 season.

The researchers selected amateur athletes of a good standard with high <u>physical performance</u> levels, who did not take part in international events, but participated in Spanish tournaments and elite cycling races.

In general, a higher percentage of professional endurance athletes expose themselves to hypoxia to aid physical improvement - 84% of professionals as opposed to 19% of amateurs, Álvarez-Herms explains.

The questionnaires show that all the athletes go through hypoxic training



because they are confident that it improves their performance. However, professionals are less optimistic than amateurs in this regard.

"Higher-level athletes believed their performances would improve by between 5% and 9%, whilst <u>amateur athletes</u> anticipated an improvement of between 10% and 48%," stated Álvarez-Herms. A possible explanation for this is that the scope for physical improvement in professional athletes is smaller than in amateurs, and therefore they are not as capable of improving.

The authors indicate that tailoring training and the use of hypoxia to each individual is key, stating: "The main recommendation we would make to sportspeople undergoing altitude <u>training</u> is to evaluate their individual physiological and physical response to altitude."

According to the expert, the conclusions drawn from the study show "a significant difference in scientific application and professional monitoring between amateur and <u>professional athletes</u> with regard to the hypoxic technique. This leads to an even greater gulf in performance between <u>athletes</u>."

**More information:** "Popularity of hypoxic training methods for endurance-based professional and amateur athletes". *Physiology* &*Behavior*. DOI: 10.1016/j.physbeh.2015.02.020

## Provided by Plataforma SINC

Citation: The majority of amateur athletes undergoing hypoxic training are not advised by specialists (2015, June 26) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2015-06-majority-amateur-athletes-hypoxic-specialists.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.