

Researchers study how marijuana impacts driving

June 24 2015



A new study conducted at the University of Iowa's National Advanced Driving Simulator has found drivers who use alcohol and marijuana together weave more on a virtual roadway than drivers who use either substance independently. However, the cocktail of alcohol and marijuana does not double the effect of the impairment.

"What we saw was an additive effect, not a synergistic effect, when we put them together," said Tim Brown, associate research scientist at NADS and corresponding author of the study. "You get what you expect if you take alcohol and [cannabis](#) and merge them together."

The study, which appears June 23 in *Drug and Alcohol Dependence*, also found participants who consumed only alcohol weaved more during a

35- to 45-minute simulated driving test than those who consumed only vaporized cannabis.

The results are part of a larger study – the first of its kind to analyze the effects of inhaled cannabis on driving performance – sponsored by the National Highway Traffic Safety Administration, National Institute of Drug Abuse and the Office of National Drug Control Policy. The UI was selected for the research because of the NADS' authenticity to real driving and the university's expertise in medicine, pharmacy, and engineering. The NADS is the only simulator of its kind that is publicly owned.

The outcomes could help shape future legislation in the United States where some experts say policies on drugged driving are woefully behind.

To date, medical [marijuana](#) is legal in 23 states and the District of Columbia while marijuana has been approved for recreational use in four states and D.C. Since legalizing [medical marijuana](#), Colorado has reported an increase in driving under the influence of cannabis cases and fatal motor vehicle crashes with cannabis-only positive drivers while states without legalized marijuana have experienced no significant change in cannabis-related crashes.

Last year, the National Highway Traffic Safety Administration's Roadside Survey of Alcohol and Drug Use by Drivers found the number of drivers with alcohol in their system has declined by nearly one-third since 2007. However, that same survey found the number of weekend nighttime drivers with evidence of drugs in their system climbed from 16.3 percent in 2007 to 20 percent in 2014. The number of drivers with marijuana in their system grew by nearly 50 percent.

Brown said plenty of research has been done on the effects of drinking alcohol and driving, but little has been done to measure the effects of

using marijuana and driving.

"Alcohol is the most common drug present in the system in roadside stops by police; cannabis is the next most common, and cannabis is often paired with alcohol below the legal limits," Brown said. "So the questions are: 'is alcohol an issue? Is cannabis an issue?' We know alcohol is an issue, but is cannabis an issue or is cannabis an issue when paired with alcohol? We tried to find out."

Researchers selected 18 participants – 13 men and five women – between the ages of 21 and 37 who reported [drinking alcohol](#) and using marijuana no more than three times a week. After spending the night at the University Hospital and Clinics to ensure sobriety, participants arrived at NADS for six "dosing visits."

First, participants were given 10 minutes to drink a mixed drink with alcohol or plain juice in an alcohol-rimmed glass and topped with alcohol to mimic alcohol taste and odor. The idea was to get the participants blood alcohol level to about .065 percent at the start of the simulated drive.

Next, they were given 10 minutes to inhale a placebo or vaporized cannabis using a vaporizing system designed in Germany called "Volcano Medic."

Once in the simulator – a 1996 Malibu sedan mounted in a 24-foot diameter dome - the [drivers](#) were assessed on weaving within the lane, how often the car left the lane and the speed of the weaving. Drivers with only alcohol in their systems showed impairment in all three areas while those strictly under the influence of vaporized cannabis only demonstrated problems weaving within the lane.

Drivers with blood concentrations of 13.1 ug/L THC, or

delta-9-tetrahydrocannabinol, the active ingredient in marijuana, showed increase weaving that was similar to those with a .08 breath alcohol concentration, the legal limit in most states. The legal limit for THC in Washington and Colorado is 5 ug/L, the same amount other states have considered.

The study also found that analyzing a driver's oral fluids can detect recent use of marijuana but is not a reliable measure of impairment.

"Everyone wants a breathalyzer which works for alcohol because [alcohol](#) is metabolized in the lungs," said Andrew Spurgin, a pharmacy research fellow on the project and co-author. "But for cannabis this isn't as simple due to THC's metabolic and chemical properties."

More information: "Cannabis Effects on Driving Lateral Control With and Without Alcohol." DOI: [dx.doi.org/10.1016/j.drugalcdep.2015.06.015](https://doi.org/10.1016/j.drugalcdep.2015.06.015)

Provided by University of Iowa

Citation: Researchers study how marijuana impacts driving (2015, June 24) retrieved 20 March 2024 from <https://medicalxpress.com/news/2015-06-marijuana-impacts.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--