

# Methadone linked to initial QTc prolongation in chronic pain

June 26 2015

---



(HealthDay)—For patients with chronic pain, methadone is associated with a small, but nonsignificant, initial increase in QTc, which does not persist, according to a study published in the June issue of *Pain Medicine*.

Samuel Grodofsky, M.D., from the University of Pennsylvania in Philadelphia, and colleagues conducted a [prospective cohort study](#) in a [chronic pain](#) clinic involving 82 patients receiving methadone and 102 receiving non-methadone opioid therapy. Automated QTc calculations from 12-lead electrocardiograms were analyzed at baseline and during the subsequent six months.

The researchers found that there was no overall higher frequency of QTc

>470 milliseconds in the methadone group (6 percent for methadone group versus 5 percent for controls;  $P = 0.722$ ). Furthermore, the methadone group had no increase in the incidence of QTc >60 milliseconds from baseline (4 and 4 percent, respectively;  $P = 0.94$ ). Patients in the methadone group exhibited an increase in QTc compared with controls in the first month after initiating methadone (5 versus 0 percent;  $P = 0.073$ ), but the difference did not persist at three and six months.

"We believe larger scale studies to further characterize the safety profile of low-dose [methadone](#) are warranted," the authors write.

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Methadone linked to initial QTc prolongation in chronic pain (2015, June 26) retrieved 26 April 2024 from  
<https://medicalxpress.com/news/2015-06-methadone-linked-qtc-prolongation-chronic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---