

Neurotoxin found in commercial seafood

June 4 2015

Popular commercial seafood purchased from Swedish supermarkets at the Stockholm region contains Beta-Methylamino-L-Alanine (BMAA), shows a doctoral thesis from Stockholm University. BMAA is a naturally-occurring amino acid with a possible link to neurodegenerative diseases such as Parkinson's disease, Alzheimer's disease and Amyotrophic lateral sclerosis. It is the first screening study to measure BMAA in commercial seafood from metropolitan markets.

"We detected BMAA in blue mussel, oyster, shrimp, plaice, char and herring but not in salmon, cod, perch and crayfish," says Liying Jiang, PhD student at the Department of Environmental Science and Analytical Chemistry (ACES), Stockholm University.

Amino acids are the building blocks of proteins. There are 20 [amino acids](#) commonly used in making proteins but BMAA is not one of them. It is not clear how BMAA should be able to lead to the onset of [neurodegenerative diseases](#). It has been discussed that BMAA may be mistakenly incorporated into proteins during synthesis and, in turn, goes on to change their normal shape and function leading to the onset or the speeding up the progress of neurodegenerative diseases.

The amino acid BMAA was first isolated and identified in extracts from the seeds of cycad trees in the West Pacific island of Guam in 1967. BMAA was potentially linked to the neurodegenerative diseases that were prevalent in Guam during the 1950's, as an important environmental risk factor.

"The BMAA producers in nature were reported to be phytoplankton, found at the bottom of the food chain. In this scenario, the entry of BMAA into the food web is inevitable." says Liying Jiang, PhD student at the Department of Environmental Science and Analytical Chemistry (ACES), Stockholm University.

According to the Food and Agricultural Organisation of the United Nations, the average consumption of seafood per capita in Sweden during 2005–2007 was estimated at 28.7 kg/year, i.e. 0.6 kg/week. Liying Jiang points out that consumers can rest assured knowing that the levels of BMAA she and her colleagues measured in commercial seafood are too low to have any direct toxic effects on animals and let alone in humans.

However, Liying Jiang emphasises that we should exercise caution due to the limited data available on the distribution and function of BMAA.

"It has been suggested that BMAA may be a slow toxin, possibly with a biomagnification potential, meaning that its levels could rise the higher up in the food chain it is found. Consequently it is important to monitor its distribution in our environment and understand the means by which it gets stored and released by organisms. These are necessary aspects that need to be addressed," says Liying Jiang, PhD student at the Department of Environmental Science and Analytical Chemistry (ACES), Stockholm University.

More information: The article is online:
www.nature.com/srep/2014/14110...1/full/srep06931.htm

Provided by Stockholm University

Citation: Neurotoxin found in commercial seafood (2015, June 4) retrieved 25 April 2024 from <https://medicalxpress.com/news/2015-06-neurotoxin-commercial-seafood.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.