

Use of osteoporosis drugs have dropped following media reports of safety concerns

June 15 2015

Following a decade of steady growth, use of bisphosphonates—medications that are effective for treating osteoporosis—declined in the United States by more than 50% from 2008 to 2012.

The sudden drop seemed to occur after media reports highlighted safety concerns, such as the development of certain fractures that occurred rarely in long-term users, despite the fact that the US Food and Drug Administration and the American Society of Bone and Mineral Research did not recommend any specific safety restrictions on <u>bisphosphonates</u>. The findings are published in the *Journal of Bone and Mineral Research*.

"Our study documents a dramatic drop in <u>osteoporosis</u> medication use by patients. Whether it will impact the long-term bone health of the nation remains to be seen," said senior author Dr. Timothy Bhattacharyya.

More information: *Journal of Bone and Mineral Research*, <u>DOI:</u> <u>10.1002/jbmr.2565</u>

Provided by Wiley

Citation: Use of osteoporosis drugs have dropped following media reports of safety concerns (2015, June 15) retrieved 25 April 2024 from <u>https://medicalxpress.com/news/2015-06-osteoporosis-drugs-media-safety.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.