

Overall rate of traumatic spinal cord injury remains stable in US

June 9 2015

Between 1993 and 2012, the incidence rate of acute traumatic spinal cord injury remained relatively stable in the U.S., although there was an increase among older adults, mostly associated with an increase in falls, according to a study in the June 9 issue of *JAMA*, a theme issue on the Americans with Disabilities Act.

Traumatic [spinal cord](#) injury leads to chronic impairment and disability. Despite the substantial effects of this injury on health-related quality of life and [health care spending](#), contemporary data on trends in incidence, causes, and medical care are limited, according to background information in the article.

Nitin B. Jain, M.D., M.S.P.H., of the Vanderbilt University School of Medicine, Nashville, Tenn., and colleagues analyzed survey data from the U.S. Nationwide Inpatient Sample (NIS) databases for 1993-2012 to examine trends in incidence, causes, [health care](#) utilization, and mortality for acute traumatic spinal cord injury.

The total study sample consisted of 63,109 patients with acute traumatic spinal cord injury. The actual number of cases in the NIS database increased from 2,659 in 1993 to 3,393 in 2012. The incidence rate for acute traumatic spinal cord injury remained relatively stable: the estimated rate was 53 cases per 1 million persons in 1993 and 54 cases per 1 million persons in 2012.

Incidence rates among the younger male population declined. For both

the male and female populations, a high rate of increase in spinal cord injury incidence from 1993 to 2012 was observed in elderly persons. Although overall in-hospital mortality increased from 6.6 percent in 1993-1996 to 7.5 percent in 2010-2012, mortality decreased significantly from 24 percent in 1993-1996 to 20 percent in 2010-2012 among persons 85 years or older.

The percentage of spinal cord injury associated with falls increased significantly from 28 percent in 1997-2000 to 66 percent in 2010-2012 in those 65 years or older. "This is a major public health issue and it likely represents a more active 65- to 84-year-old U.S. population currently compared with the 1990s, which increases the risk of falls in this age group. This issue may be further compounded in the future because of the aging population in the United States," the authors write.

More information: *JAMA*, [DOI: 10.1001/jama.2015.6250](https://doi.org/10.1001/jama.2015.6250)

Provided by The JAMA Network Journals

Citation: Overall rate of traumatic spinal cord injury remains stable in US (2015, June 9)
retrieved 10 April 2024 from

<https://medicalxpress.com/news/2015-06-traumatic-spinal-cord-injury-stable.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.