


Viral commuters: How influenza viruses use transportation systems in the US

June 18 2015



This photo of the US Interstate Commuter Network shows the number of people traveling daily between states for work. Credit: Bozick, CC-BY

In increasingly mobile modern societies, long-distance transmission can rapidly spread pathogens. A study published on June 18th in *PLOS Pathogens* suggests that both airline and commuter road travel influence flu virus distribution in the continental U.S.

When viruses invade naïve host populations and are propagated predominantly by local transmission, we expect to observe wave-like spread across geographic space. As viruses evolve rapidly, because of their high mutation rate, these wave-like patterns of local transmission (from person-to-person and village-to-village) should generate wave-like patterns of [genetic variation](#) where the geographic distance between locations and the genetic distance between variants is positively correlated.

In today's world, however, transmission patterns are more complicated, as human pathogens also 'travel' by road, rail, and air. To examine how genetic variation correlates with spatial distribution in a highly mobile society, Brooke Bozick and Leslie Real, from Emory University in Atlanta, explored whether measures of distance defined by airline and commuter transportation networks can explain the population genetic structure of [seasonal influenza](#) viruses within the U.S.

Analyzing the travel networks, the researchers calculated that during the flu season, approximately 1.6 million people travel along the interstate aviation network per day, and that most U.S. states are well connected to most other states. More people (over 3.8 million) travel daily across the interstate ground travel commuter network, but the vast majority of connections here occur between neighboring states, and more so in the Eastern than in the Western U.S.

Next the researchers analyzed the genetic variation of flu viruses (both the H3N2 and the H1N1 viruses) from viruses isolated and classified over 10 seasons: from 2003/2004 to 2012/2013 across the U.S. They then compared genetic distance (which is greater between sequences that are more divergent) with (1) geographic distance, and with the alternative distance measures of (2) interstate airline connections and (3) interstate commuter connections.

They found that while all three distance metrics were correlated with genetic distance at least some of the time, the commuter network was more often associated with genetic distance than any other measure of spatial or network distance for the H1N1 subtype. Further, the association between genetic distance and the commuter network often remained significant after geographic distance is taken into account, demonstrating that the relative magnitude of host movement over space has a greater influence on the route of pathogen spread than the geographic proximity of sampling locations.

The researchers say that to their knowledge, this is the first evidence that population structure (as revealed by travel networks) for seasonal influenza is detectable at the scale of the continental U.S. In contrast, population structure was not detected in most of the data analyzed for H3N2 virus, even though both geographic distance and commuter distance were, at times, correlated with [genetic distance](#). It might be that the current resolution of the origin of the virus variants (by state rather than by county) is not sufficient to detect a correlation between the usually more aggressive and fast-spreading H2N3 virus and travel patterns.

The researchers conclude that while recent findings have shown that the aviation network plays an important role in the world-wide transmission of seasonal influenza, their results suggest that when [population structure](#) is detectable, it is the commuter network that is of greater importance at more regional scales. Discussing the public health implications, the researchers say that the detection of network structure implies that patterns of epidemic spread are, to some extent, predictable and point out that 'the absence of predictability is problematic for the design of containment strategies, since it suggests that the annual seasonal spread of influenza within countries is highly variable and depends heavily on chance events.'

The findings here, based on genetic data, are in agreement with previous research using influenza-like illness case data; both show that patterns do exist, with strongly connected states having similarly timed epidemic peaks and similar genetic variants circulating. The patterns they found are likely influenced by states with many commuters, and the researchers suggest that 'the identification of these states, as well as network pathways that contribute substantially to influenza spread, is an important next step for epidemiological research.'

More information: *PLOS Pathogens* ,
[dx.plos.org/10.1371/journal.ppat.1004898](https://doi.org/10.1371/journal.ppat.1004898)

Provided by Public Library of Science

Citation: Viral commuters: How influenza viruses use transportation systems in the US (2015, June 18) retrieved 20 March 2024 from <https://medicalxpress.com/news/2015-06-viral-commuters-influenza-viruses.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
