

## West African Ebola virus strain less virulent than prototype 1976 strain

June 9 2015



The Ebola virus, isolated in November 2014 from patient blood samples obtained in Mali. The virus was isolated on Vero cells in a BSL-4 suite at Rocky Mountain Laboratories. Credit: NIAID

What:The Makona strain of Ebola virus (EBOV) circulating in West Africa for the past year takes roughly two days longer to cause terminal



disease in an animal model compared to the original 1976 Mayinga strain isolated in Central Africa, according to a new National Institutes of Health (NIH) report. The results provide important information to scientists who have wondered if the Ebola virus in West Africa is becoming more severe. In fact, the new study suggests the current virus has a decreased ability to cause disease in their animal model compared to the 1976 strain.

Using cynomolgus macaques to model infection and disease in humans, scientists from NIH's National Institute of Allergy and Infectious Diseases (NIAID) infected three animals with 1976 EBOV-Mayinga and three with 2014 EBOV-Makona. While both groups of animals were shedding virus three days after being infected, those in the Mayinga group developed a rash on day four and became extremely ill at days five and six. Those in the Makona group did not develop a rash until six days after infection, and severe disease appeared on days seven and eight. Further, liver damage—typical in Ebola disease—was delayed by about two days in the Makona group compared to the Mayinga group.

Of note, the scientists reported that the immune system of animals in the Makona group produced about three times the amount of a virus-fighting protein, interferon gamma, compared to the Mayinga group. They plan more studies of the immune response, but believe that at least seven days are needed after Ebola infection to mount an effective response. This response does not seem to develop during EBOV-Mayinga infection in cynomolgus macaques because disease progression is too fast.

**More information:** A Marzi et al. Delayed disease progression in cynomolgus macaques infected with Ebola virus Makona strain. *Emerging Infectious Diseases* DOI: 10.3201/eid2110.150259 (2015).



## Provided by NIH/National Institute of Allergy and Infectious Diseases

Citation: West African Ebola virus strain less virulent than prototype 1976 strain (2015, June 9) retrieved 2 May 2024 from https://medicalxpress.com/news/2015-06-west-african-ebola-virus-strain.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.