


# Elective surgery is associated with lower risk of death than drugs for ulcerative colitis

July 13 2015

---



Endoscopic image of a bowel section known as the sigmoid colon afflicted with ulcerative colitis. The internal surface of the colon is blotchy and broken in places. Credit: Samir/Wikipedia

Patients over 50 with ulcerative colitis (UC), a chronic disease of the colon, who undergo surgery to treat their condition live longer than those who are treated with medications, according to a new study from the Perelman School of Medicine at the University of Pennsylvania. The results are published this week in *Annals of Internal Medicine*.

"Ulcerative colitis is a chronic [disease](#) that most physicians opt to treat with medications, as opposed to [surgery](#)," said the study's lead author Meenakshi Bewtra, MD, PhD, MPH, assistant professor of Medicine and Epidemiology at Penn. "This new finding highlights a potential unrecognized advantage of a surgical approach to the disease."

Ulcerative colitis is a type of [inflammatory bowel disease](#) characterized by inflammation in the colon, or the [large intestine](#). The inflammation can cause abdominal discomfort, bleeding, and diarrhea. It affects as many as 700,000 Americans, according to the Crohn's & Colitis Foundation of America. Men and women are equally affected, with most [patients](#) diagnosed by their mid-30s.

The convention in treatment of UC is that surgery is considered a treatment of last resort. However, [medical](#) treatments, involving immunosuppressant and steroid drugs, come with significant side effects, and can increase the risk of infection and some cancers. These medications effectively control the disease in less than 50 percent of patients.

In a retrospective study, Bewtra and colleagues analyzed data from the Centers for Medicare and Medicaid Services (CMS) to examine whether patients with advanced UC pursuing elective colectomy - in which surgeons remove the patient's colon—had improved survival compared to similar patients pursuing chronic drug therapy. The researchers defined advanced UC as those patients who had had at least one hospitalization for UC, had two or more corticosteroid prescriptions

within a 90-day period, or had any prescription for immunosuppressant therapy.

The study identified 32,833 UC patients 18 years of age and older who had at least six months of Medicare/Medicaid eligibility and who fit the inclusion criteria. Of these, 830 underwent elective colectomy and the remaining 32,003 were managed on drug therapies.

Mortality rates associated with elective colectomy and medical therapy were 34 and 54 per 1,000 person-years, respectively, thus showing that elective colectomy was associated with an improved survival rate. A person-year is a measure of survival that is equivalent to the number of years a person is followed multiplied by the number of subjects in the study. In further analysis, the survival benefit seemed greatest in patients age 50 and over. The findings also suggested a benefit for surgery in those patients who could not sustain a disease remission - defined as stable disease without the need for hospitalization or changes to drug therapy—on immunosuppressant therapy, but further work is needed to confirm these results.

"With this new knowledge, physicians should be empowered to begin a dialogue about surgery earlier in their patients' course of treatment," Bewtra said. "Many patients are afraid of surgical therapy for UC. This study should help them to understand that the benefits of surgery may extend beyond just reducing the symptoms of uncontrolled UC."

Provided by University of Pennsylvania School of Medicine

Citation: Elective surgery is associated with lower risk of death than drugs for ulcerative colitis (2015, July 13) retrieved 6 May 2024 from <https://medicalxpress.com/news/2015-07-elective-surgery-death-drugs-ulcerative.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.