


# Study identifies factors affecting prescription pain reliever misuse

July 10 2015



Researchers at the UGA School of Social Work found that adults aged 50 and above were more likely to acquire prescription pain relievers through multiple doctors, whereas younger individuals were more likely to acquire them from

friends, relatives, or drug dealers. Regardless of age, all those who misused prescription painkillers had a history of recent use of illicit drugs. Credit: UGA

People who misuse prescription pain relievers all have one thing in common, University of Georgia researchers have discovered: a history of recent illicit drug use. How they acquire such drugs varies according to age, however. The findings, published in the journal *Addictive Behaviors*, may help health care providers and others curb painkiller misuse.

In a nationwide study, researchers from the UGA School of Social Work found that individuals of any age who used illicit drugs such as marijuana, cocaine or heroin within the past year had a higher likelihood of misusing prescription pain relievers as well. A study just released by the Centers for Disease Control, which found that heroin use was highest among those who abused cocaine or opioid pain relievers within the past year, buttressed the UGA research.

"Male or female, black or white, rich or poor, the singular thing we found was that if they were an illicit drug user, they also had many, many times higher odds of misusing [prescription pain relievers](#)," said Orion Mowbray, an assistant professor in the School of Social Work and the UGA study's lead author.

The UGA researchers also determined that adults aged 50 and above were more likely to acquire pain relievers through more than one doctor, whereas younger individuals were more likely to acquire them from friends, relatives or drug dealers.

The investigation was based on more than 13,000 responses to the National Survey on Drug Use and Health. The annual survey, sponsored

by the U.S. Department of Health and Human Services, collects data on the use of tobacco, alcohol, [illicit drugs](#), [prescription drugs](#), and mental health problems among individuals aged 12 and older.

Prescription pain relievers represent the majority of all prescription drugs that are misappropriated in the U.S., and misuse has risen dramatically in recent years. The most over-used pain relievers are opioids—addictive painkillers such as codeine, oxycodone and morphine.

According to a 2013 report by the Substance Abuse and Mental Health Services Administration, a unit of the HHS, emergency room treatments for opioid misuse—including suicide attempts and accidental overdoses—increased 183 percent from 2004 to 2011.

The reason for the increase is not clear, but the report offers possible solutions to address the problem.

"If we know how people come to possess the [pain relievers](#) they misuse, we can design better ways to lower that likelihood," said Mowbray. "This study gives us the knowledge we need to substantially reduce the opportunities for misuse."

The study calls for greater coordination between medical care providers to reduce the possibility of over-prescription of painkillers, and for improving the communication between doctors, patients and the public.

"Doctors may conduct higher quality conversations with older patients about the consequences of drug use before they make any prescription decisions, while families and friends should know about the substantial health risks before they supply a young person with a prescription pain reliever," said Mowbray.

Adam Quinn, a doctoral student in the School of Social Work, contributed to the research. The study, "Prescription pain reliever misuse prevalence, correlates, and origin of possession throughout the life course," is available online at <http://www.sciencedirect.com/science/article/pii/S030646031500194X>.

Provided by University of Georgia

Citation: Study identifies factors affecting prescription pain reliever misuse (2015, July 10) retrieved 2 May 2024 from <https://medicalxpress.com/news/2015-07-factors-affecting-prescription-pain-reliever.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--