

## **Repeat infection with malaria parasites might make mosquitoes more dangerous**

July 16 2015



If they have the opportunity mosquitoes will blood feed every 2-5 days and therefore can be exposed to multiple infections. Credit: Sarah Reece & Sinclair Stammers, CC-BY

In malaria-endemic regions, humans are often infected repeatedly with



the *Plasmodium* parasite, and the consequences of such multiple infections are under intense study. In contrast, little is known about possible co-infection and its consequences in the *Anopheles* mosquitoes that transmit the disease. A study published on July 16th in *PLOS Pathogens* reports that not only can individual mosquitoes accumulate infections from multiple blood feeds, but also that an existing malaria infection makes mosquitoes more susceptible to a second infection, and that infections reach higher densities when another strain is already present.

Interested in interactions between malaria parasites and their insect hosts, Laura Pollitt, from the University of Edinburgh, UK, and colleagues in the US, asked whether and how mosquitoes can be infected with different *Plasmodium* strains, how such heterogeneous parasites interact in the insects, and whether such interactions affect transmission of malaria to vertebrate hosts.

The researchers set up cages of female *Anopheles* mosquitoes and allowed them at defined times to feed on mice infected with two different *Plasmodium* strains. This study design allowed them to examine how the presence of a co-infecting strain affects parasites that enter the vector first and second, and to test whether co-<u>infection</u> impacts mosquito survival.

They found that mosquitoes can accumulate mixed strain malaria infections after feeding on multiple hosts, and found that parasites have a greater chance of establishing a secondary infection if another *Plasmodium* strain is already present in a mosquito. Moreover, the presence of the primary infection facilitated replication of the secondary infection while the first infection developed as normal. This resulted in doubly infected mosquitoes having substantially higher parasite loads.

The large parasite numbers do not appear to kill the insects, and as it is



expected that mosquitoes carrying more parasites are more likely to transmit malaria to vertebrates, mosquitoes taking multiple infective bites might disproportionally contribute to malaria transmission. This in turn would increase rates of mixed infections in vertebrate (including human) hosts, with implications for the evolution of parasite virulence and the spread of drug-resistant strains.

"If the facilitation we have demonstrated here", the authors say, "occurs in natural transmission settings to humans, there could be significant epidemiological consequences. Control measures reducing prevalence in the vertebrate host, and therefore reducing the likelihood of mosquitoes taking multiple infective feeds, could disproportionally reduce transmission of individual strains. By increasing the proportion of infectious <u>mosquitoes</u> with mixed strain infections, it is also likely that the facilitation reported here will increase the rates of mixed infections in vertebrate hosts which could have implications for infection virulence and the spread of drug resistant strains."

**More information:** Pollitt LC, Bram JT, Blanford S, Jones MJ, Read AF (2015) Existing Infection Facilitates Establishment and Density of Malaria Parasites in Their Mosquito Vector. PLoS Pathog 11(7):e1005003. DOI: 10.1371/journal.ppat.1005003

Provided by Public Library of Science

Citation: Repeat infection with malaria parasites might make mosquitoes more dangerous (2015, July 16) retrieved 27 April 2024 from <u>https://medicalxpress.com/news/2015-07-infection-malaria-parasites-mosquitoes-dangerous.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.