

Research letter: Indoor tanning rates drop among US adults

July 1 2015

Indoor tanning rates dropped among adults from 5.5 percent in 2010 to 4.2 percent in 2013, although an estimated 7.8 million women and 1.9 million men still engage in the practice, which has been linked to increased cancer risk, according to the results of a study published online in a research letter by *JAMA Dermatology*.

Gery P. Guy Jr., Ph.D., M.P.H., of the Centers for Disease Control and Prevention, Atlanta, and coauthors analyzed data for 59,145 individuals from the 2010 and 2013 National Health Interview Survey, a nationally representative sample of U.S. adults.

In addition to the overall reduction, the authors identified decreases in indoor tanning rates among individuals ages 18 to 29 (11.3 percent in 2010 to 8.6 percent in 2013), women (8.6 percent in 2010 to 6.5 percent in 2013), and men (2.2 percent in 2010 to 1.7 percent in 2013).

Among women who indoor tanned, the frequency was 28 percent lower among the oldest group, 45 percent lower among college graduates, 33 percent lower among women in fair or poor health and 23 percent lower among [women](#) meeting aerobic or strength physical activity criteria. However, indoor tanning frequency among men was 177 percent higher among men age 40 to 49 and 71 percent higher in [men](#) age 50 or older but 45 percent lower among cancer survivors, according to the results.

The authors suggest the decrease in indoor tanning may be partly attributable to increased awareness of its harms as indoor tanning

devices have been classified as carcinogenic to humans, laws restricting access by minors have been enacted and a 10 percent excise tax on indoor tanning has been implemented.

The authors caution a causal inference cannot be made between behaviors and the frequency of indoor tanning from their data.

"Research regarding the motivations of indoor tanners could inform the development of new interventions. Physicians can also play a role through behavioral counseling, which is recommended for fair-skinned persons aged 10 to 24 years. Continued surveillance of [indoor tanning](#) will aid program planning and evaluation by measuring the effect of [skin cancer prevention](#) policies and monitoring progress," the study concludes.

More information: *JAMA Dermatology*. Published online July 1, 2015. [DOI: 10.1001/jamadermatol.2015.1568](https://doi.org/10.1001/jamadermatol.2015.1568)

Provided by The JAMA Network Journals

Citation: Research letter: Indoor tanning rates drop among US adults (2015, July 1) retrieved 25 April 2024 from <https://medicalxpress.com/news/2015-07-letter-indoor-tanning-adults.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--