


No lasting value for minimally invasive lumbar laminotomy

July 29 2015



(HealthDay)—For patients with neural foraminal or lateral recess stenosis with unilateral leg neurogenic symptoms (NS), a minimally invasive surgical (MIS) approach offers no advantage over an open lumbar laminotomy approach in the longer term, according to a study published in the Aug. 1 issue of *The Spine Journal*.

Chia-Liang Ang, M.B.B.S., M.Med., from the Singapore General Hospital, and colleagues compared postoperative improvements in functional and pain scores for open versus MIS lumbar laminotomy in a retrospective study. Data were reviewed for 113 patients who underwent MIS (83 patients) or open (30 patients) unilateral one-level lumbar laminotomy for treatment of neural foraminal or lateral recess stenosis with unilateral leg NS.

The researchers observed no differences in improvement in back or [leg pain](#), or in improvement in Oswestry Disability Index, NS, or Short Form-36 scores at six or 24 months after surgery. At six, but not 24, months there was greater satisfaction with treatment in the MIS group ($P = 0.009$). Within 24 months, three [patients](#) experienced an inadvertent durotomy and two underwent fusion of the operated segment in the MIS group.

"Compared with an open approach, MIS lumbar laminotomy gave no clear advantages in longer-term functional or [pain scores](#)," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: No lasting value for minimally invasive lumbar laminotomy (2015, July 29) retrieved 2 May 2024 from <https://medicalxpress.com/news/2015-07-minimally-invasive-lumbar-laminotomy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
