

Failed communication associated with readmission

July 16 2015



(HealthDay)—Failed communication attempts are associated with readmission among Medicare beneficiaries with congestive heart failure, although the correlation is no longer significant after adjustment for other variables, according to a study published online June 26 in the *Journal of the American Geriatrics Society*.

Matthew J. Press, M.D., from the Weill Cornell Medical College in New York City, and colleagues conducted a retrospective cohort study involving 5,698 Medicare beneficiaries with <u>congestive heart failure</u> who received home health care from the Visiting Nurse Service of New York (VNSNY) after hospital discharge. The authors examined the correlation between patient-level measures of communication failure and all-cause readmission.



The researchers found observed high external validity using <u>natural</u> <u>language processing</u> to identify failed communication attempts (kappa, 0.850; P

"Natural language processing of electronic medical records can be used to identify failed communication attempts between home health nurses and physicians, but other variables mostly explained the association between <u>communication failure</u> and readmission," the authors write.

Several authors are employees of VNSNY; funding for the study was provided by the Aetna Foundation.

More information: <u>Abstract</u>

Full Text (subscription or payment may be required)

Copyright © 2015 HealthDay. All rights reserved.

Citation: Failed communication associated with readmission (2015, July 16) retrieved 28 April 2024 from <u>https://medicalxpress.com/news/2015-07-readmission.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.