

# Study reveals brain network responsible for cognitive changes in multiple sclerosis

July 7 2015

---



Bart Rypma, Ph.D., Principal Investigator, M.S. study, Center for BrainHealth.  
Credit: Randy Anderson

An estimated 2.3 million individuals are living with multiple sclerosis (MS) worldwide. Approximately half of all individuals with MS

experience changes in cognition such as impaired concentration, attention, memory, and judgment. The underlying brain basis for these deleterious effects has been largely elusive. New findings published yesterday in *Neuropsychology* reveal that decreased connectivity between network-specific brain regions are to blame for the central deficit common to the various cognitive changes associated with MS, slowed cognitive speed.

In the first study of its kind, researchers at the Center for BrainHealth at The University of Texas at Dallas and The University of Texas Southwestern Medical Center found that, compared to healthy controls, individuals with MS exhibit weaker brain connections between the dorsolateral prefrontal cortex and posterior [brain regions](#). The change amounts to a breakdown in communication between the part of the brain responsible for executing goal-directed thought and action and the regions responsible for carrying out tasks related to cognitive speed such as visual processing, motor execution, and object recognition. The researchers believe that the diminished connections are likely the result of decreased white matter surrounding the neurons in the brain.

"Our study is the first to really zero in on the physiology of cognitive speed, the central cognitive deficit in MS," explained Center for BrainHealth principal investigator Bart Rypma, Ph.D., who also holds the Meadows Foundation Chair at UT Dallas. "While white matter is essential to efficient network communication, white matter degradation is symptomatic of MS. This study really highlights how tightly coupled connectivity is to performance and illuminates the larger, emerging picture of white matter's importance in human cognitive performance."

Collaborating with Elliot Frohman, M.D., Ph.D., director of the Multiple Sclerosis Program and Clinical Center at UT Southwestern, the study recruited 29 participants with relapsing-remitting MS and 23 age- and sex- matched healthy controls. Participants underwent functional

magnetic resonance imaging (fMRI) while completing a measure of cognitive processing speed. Participants were given 4 seconds to view a nine-item key of number and symbol pairs (for example '+' above the number 3) and one number-symbol pair probe. Participants were asked to indicate with a left or right thumb button press whether or not the probe appeared in the key.

While accuracy was similar for both healthy controls and individuals with MS, response times for individuals with MS were much slower. Analysis of the fMRI data revealed that while completing this measure, MS patients showed weaker functional connections with [dorsolateral prefrontal cortex](#).

"These findings reveal a diffuse pattern of disconnectivity with executive areas of the brain," explained the study's lead author, Nicholas Hubbard, a doctoral candidate at the Center for BrainHealth working with Dr. Rypma. "Importantly, these decreases in connectivity predicted MS-related cognitive slowing both in and out of the fMRI environment suggesting that these results were not specific to our task, but rather were able to generalize to other situations where cognitive speed is required."

This research supports the need for therapies that target white matter structures and white matter proliferation. Rypma and Hubbard are currently conducting research to further explore the physiology of [white matter](#) to better understand cognitive speed reductions not only in MS, but also in healthy aging individuals.

**More information:** *Neuropsychology*,  
[psycnet.apa.org/psycinfo/2015-30063-001](http://psycnet.apa.org/psycinfo/2015-30063-001)

Provided by Center for BrainHealth

Citation: Study reveals brain network responsible for cognitive changes in multiple sclerosis (2015, July 7) retrieved 23 April 2024 from <https://medicalxpress.com/news/2015-07-reveals-brain-network-responsible-cognitive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.