

Static synapses on a moving structure: Mind the gap!

July 22 2015



Axon initial segments (blue) and the synapses that form along them (red and green) are shown. Credit: Winnie Wefelmeyer. MRC Centre for Developmental Neurobiology

In biology, stability is important. From body temperature to blood pressure and sugar levels, our body ensures that these remain within reasonable limits and do not reach potentially damaging extremes. Neurons in the brain are no different and, in fact, have developed a number of ways to stabilise their electrical activity so as to avoid becoming either overexcitable, potentially leading to epilepsy, or not excitable enough, leading to non functional neurons. A new study published in *PNAS* by researchers from the MRC Centre for Developmental Neurobiology characterises a novel way in which neurons remain electrically stable when confronted with chronic increases in neuronal activity.



The site at the centre of this control mechanism is a short segment of the axon, where <u>electrical activity</u> is initiated in the first place. Also known as the axon initial segment (or AIS for short), this remarkable structure is responsible for integrating all the information that a neuron receives via its synapses to produce an action potential, the electrical currency of information used by neurons. It is perhaps no surprise then that modulating this domain should have an important impact on a cell's excitability. In fact, previous findings from the Burrone lab were the first to uncover that changes in AIS structure could alter the excitability of neurons to stabilise their overall electrical activity (Grubb & Burrone Nature 365:1070). What is less known is what happens to the unique synapses that form along the AIS, when this region of the axon is modified. "We know very little about these odd axonal synapses, other than they are in the right place to modulate neuronal output by acting directly on the axon initial segment", says senior author Prof Juan Burrone, "but what happens when the AIS changes was, until now, a mystery".

This study shows for the first time that in slices of the hippocampus, a region of the brain known to be important for learning and memory, increases in neuronal activity causes the AIS to move away from the cell body, along the axon, by about half its length. The synapses however stay in place, causing a mismatch between the AIS and the synapses that control it. In collaboration with Dr Daniel Cattaert at the INCIA in Bordeaux, the authors used computational models to study the functional consequences of this arrangement. This approach revealed that the synapses that were left behind, those that lie in the gap between the cell body and the AIS, are particularly important for decreasing neuronal excitability, allowing neurons to remain functional even when under constant stimulation. As Dr Winnie Wefelmeyer, the leading author of this study puts it: "The axon initial segment is like the vocal chord of the neuron: without it, it would be unable to communicate. Changing the position of the AIS relative to the modulating synapses ensures that what



the neuron is saying stays meaningful".

More information: Wefelmeyer W, Cattaert D, Burrone J (2015). Activity-dependent mismatch between axo-axonic synapses and the axon initial segment controls neuronal output. *PNAS* (2015) <u>DOI:</u> 10.1073/pnas.1506823112

Provided by King's College London

Citation: Static synapses on a moving structure: Mind the gap! (2015, July 22) retrieved 2 May 2024 from <u>https://medicalxpress.com/news/2015-07-static-synapses-mind-gap.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.