

Weight loss predicts mortality in rheumatoid arthritis

July 13 2015



(HealthDay)—Weight loss is a strong predictor of mortality in patients with rheumatoid arthritis (RA), according to a study published online June 26 in *Arthritis & Rheumatology*.

Joshua F. Baker, M.D., from the Philadelphia Veterans Affairs (VA) Medical Center, and colleagues identified 1,674 RA [patients](#) from the VA RA Registry and analyzed their medical records to identify body mass index (BMI) at each study visit, as well as date of death, if applicable.

The researchers noted 312 deaths over 9,183 person-years. After adjusting for demographics, comorbidities, BMI, smoking, and RA therapies, a loss in BMI of ≥ 1 kg/m² was associated with a greater risk

of death (hazard ratio [HR], 1.99; P weight loss at an annualized rate of ≥ 3 kg/m² (HR, 2.49; P

"These observations may explain the observed obesity paradox and do not support a biologically protective role of obesity," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Weight loss predicts mortality in rheumatoid arthritis (2015, July 13) retrieved 6 May 2024 from <https://medicalxpress.com/news/2015-07-weight-loss-mortality-rheumatoid-arthritis.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|