

## Azithromycin routine in hospital despite risk for QTc prolongation

August 31 2015



(HealthDay)—Azithromycin is routinely prescribed to hospitalized patients despite risk factors for corrected QT (QTc) prolongation and administration of interacting medications, according to a study published online Aug. 19 in the *Journal of Hospital Medicine*.

Rachael A. Lee, M.D., from the University of Alabama at Birmingham, and colleagues retrospectively analyzed data from 100 adult inpatients randomly selected from 1,610 patient encounters (October 2012 to April 2013), all of whom were treated with at least one dose of azithromycin. Telemetry charges and baseline electrocardiogram (ECG) prior to azithromycin administration were evaluated.

The researchers found that 79 percent of azithromycin use was empiric. Of the 65 patients who received a baseline ECG prior to prescribing



azithromycin, 60 percent had borderline or abnormal QTc prolongation. There were more abnormal ECGs at baseline (P = 0.03) among the 76 percent of patients who were prescribed two or more QTc-prolonging medications in addition to azithromycin, despite having telemetry ordered less than half of the time.

"Selection of azithromycin by providers appears to be independent from these <u>risk factors</u>, and education and vigilance to drug-drug interactions may be useful in limiting <u>cardiac events</u> with prescribing <u>azithromycin</u>," the authors write.

**More information:** Abstract

Full Text (subscription or payment may be required)

Copyright © 2015 HealthDay. All rights reserved.

Citation: Azithromycin routine in hospital despite risk for QTc prolongation (2015, August 31) retrieved 19 April 2024 from

https://medicalxpress.com/news/2015-08-azithromycin-routine-hospital-qtc-prolongation.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.