

New biomarkers show exercise helps reduce daytime sleep disorder

August 11 2015



Dr. Madhukar Trivedi. Credit: UT Southwestern

Aerobic exercise can help alleviate excessive daytime sleepiness among depressed individuals, researchers with UT Southwestern Medical Center's Center for Depression Research and Clinical Care have found.

Researchers looking at blood samples identified two biological markers for the condition, called hypersomnia, which is characterized by sleeping too much at night as well as [excessive daytime sleepiness](#), in those with Major Depressive Disorder. Exercise reduced the levels of the two

biomarker proteins, resulting in reduced excessive sleepiness, the researchers found.

"Hypersomnia, as well as insomnia, have been linked in the development, treatment, and recurrence of [depression](#). Sleep disturbances are also some of the most persistent symptoms in depression. Identifying these biomarkers, combined with new understanding of the important role of exercise in reducing hypersomnia, have potential implications in the treatment of [major depressive disorder](#)," said senior author Dr. Madhukar Trivedi, Director of UT Southwestern's Center for Depression Research and Clinical Care, and Chief of the Mood Disorders Division of Psychiatry at UT Southwestern.

People with hypersomnia are compelled to nap repeatedly during the day, often at inappropriate times such as at work, during a meal, or in conversation. They often have difficulty waking from a long sleep, and may feel disoriented upon waking, according to the National Institute of Neurological Disorders and Stroke (NINDS), part of the National Institutes of Health (NIH). Other symptoms may include anxiety, increased irritation, decreased energy, restlessness, slow thinking, slow speech, loss of appetite, hallucinations, and memory difficulty. Some patients lose the ability to function in family, social, occupational, or other settings.

Researchers had previously found a negative loop in which sleep, inflammation and depression interact and progressively worsen. The results of the current and previous research on insomnia suggest that exercise may be resetting this negative feedback loop, said Dr. Trivedi, who holds the Betty Jo Hay Distinguished Chair in Mental Health, and received the 2015 American Psychiatric Association Award for Research, the Association's most significant award for research.

The researchers were able to identify the biomarkers based on blood samples provided by participants in the Treatment with Exercise Augmentation for Depression (TREAD) study, who were randomly assigned to two types of [aerobic exercise](#) to determine the effects of exercise on their depression. More than 100 subjects ages 18 to 70 who had Major Depression Disorder participated, and as part of the study had also agreed to provide blood samples.

In this study, researchers examined four biomarkers - brain-derived neurotrophic factor (BDNF) and inflammatory cytokines called tumor necrosis factor alpha, and two interleukins, IL-1 β and IL-6, from [blood samples](#) collected before and after the 12-week exercise intervention.

Researchers found that reductions in two biomarkers, BDNF and IL-1 β , are related to reductions in hypersomnia. "Identification of biomarkers that uniquely predict or correlate with improvements in hypersomnia and insomnia is an important step toward more effective treatment of MDD," said lead author Dr. Chad Rethorst, Assistant Professor of Psychiatry with the Center for Depression Research and Clinical Care.

Previous analysis of the TREAD data demonstrated significant reductions in insomnia symptoms with [exercise](#), but the two biomarkers identified above did not correlate to changes in insomnia, Dr. Rethorst said. They did find, however, that lower baseline levels of IL-1 β were predictive of greater improvements in insomnia. The findings suggest distinct mechanisms are involved in insomnia versus hypersomnia, and that further research will be needed to identify the appropriate [biomarkers](#) for [insomnia](#).

More information: [www.nature.com/tp/journal/v5/n ... full/tp2015104a.html](http://www.nature.com/tp/journal/v5/n...full/tp2015104a.html)

Provided by UT Southwestern Medical Center

Citation: New biomarkers show exercise helps reduce daytime sleep disorder (2015, August 11)
retrieved 4 May 2024 from

<https://medicalxpress.com/news/2015-08-biomarkers-daytime-disorder.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.