

Surprise cardiac finding predicts future risk

August 30 2015

In patients with chronic ischemic heart disease, a small left ventricle with thick walls, is the strongest predictor of morphologic remodelling, which is generally considered a first step towards heart failure, according to unexpected findings presented today at ESC Congress 2015.

Results of the DOPPLER-CIP (which stands for "Determining Optimal non-invasive Parameters for the Prediction of Left vEntricular morphologic and functional Remodeling in Chronic Ischemic Patients") study were not expected and, if confirmed by other studies, "could completely change risk stratification among patients with stable [coronary artery disease](#)," according to the study coordinators Frank Rademakers, MD and Jan D'Hooge, PhD.

"We were indeed surprised by these findings," said the investigators, who are from the University of Leuven, Belgium. "The general belief is that larger ventricles with thin walls (a typical 'infarct ventricle') would be at higher risk of remodelling, with a possible explanation for this being that there is increased wall stress in such hearts. But our findings show that it is actually small hearts with thick walls that are more at risk. As this goes against general belief, we have checked and re-checked our data, and analysis, and have run several consistency tests, but they all led to this same conclusion."

There are currently no guidelines for assessing a patient's risk for this type of deterioration, they noted.

DOPPLER-CIP compared different non-invasive methods to determine

the most useful tool at baseline for predicting risk of cardiac remodelling two years later.

The study included 676 patients, from 6 European countries, with suspicion of chronic [ischemic heart disease](#).

The patients underwent standard diagnostic tests at baseline including: electrocardiogram (ECG), exercise testing with continuous ECG monitoring, and measurement of [maximal oxygen uptake](#) (VO₂max), as well as blood sampling and quality of life assessments.

In addition to these standard tests, patients also underwent at least two stress imaging tests including echocardiography (ECHO), magnetic resonance imaging (MRI) and/or single positron emission computed tomography (SPECT) stress test, stress ECHO and stress MRI.

After these baseline evaluations all [patients](#) received optimal, guideline-based treatment including revascularization, partial revascularization, or pharmacologic treatment at their physician's discretion.

At the end of the study period, about 20% of the subjects had evidence of cardiac remodelling based on MRI or ECHO results, with the best baseline predictors of this remodelling being [left ventricular](#) size measured as the "left ventricular end-diastolic volume" (LV EDV) and left ventricular mass (LVM) .

Specifically, a small LV end-diastolic volume (< 145 ml) at baseline had a 25-40% chance of remodelling, compared to a larger EDV which had a decreased risk (20%; P

Citation: Surprise cardiac finding predicts future risk (2015, August 30) retrieved 4 May 2024 from <https://medicalxpress.com/news/2015-08-cardiac-future.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.