


# **Chickenpox vaccination does increase shingles cases, but mainly in young adults**

August 11 2015

---

## Benefits of re-exposure to children with chicken pox shown in years

- Common belief of lasting benefits or re-exposure to children with chicken pox (based on model-based prediction).
- Actual lasting benefits of re-exposure to children with chicken pox based on new model that incorporates biological data from individuals.



Re-exposure to chickenpox as adult, having experienced it as a child, boosts immunity to shingles for only two years. This contrasts to the 20 years previously thought. Credit: *eLife*

Vaccinating one-year-olds against chickenpox could temporarily nearly double the incidence of shingles in the wider population, but in younger adults than previously thought.

The effect occurs because vaccination reduces the likelihood of adults who experienced [chickenpox](#) as a child being re-exposed to the virus. Re-exposure boosts immunity to shingles, caused by the same virus, Varicella-zoster virus.

In a study to be published in the journal *eLife*, scientists from the Universities of Antwerp and Hasselt (Belgium) have predicted that the temporary effect of a rise in shingles cases dominates in 31 to 40-year-olds. This is younger than previously predicted and this age group is less at risk of developing the most serious shingles symptoms. Many countries have avoided introducing universal chickenpox vaccination in children because it was previously predicted that the reduction in chickenpox related disease would be outbalanced by the temporarily increase in shingles-related disease.

A new model developed by the scientists also confounds previous findings on the length of time re-exposure chickenpox boosts immunity to shingles. The effect was thought to last for up to 20 years, but results of the current modeling study show it only lasts for two. The new model is the first based on real immunological and virological data from individuals.

"We were surprised to find that re-exposure to chickenpox is beneficial for so few years and also that the most pronounced effect of vaccination on increasing cases of shingles is in [younger adults](#)," says lead author Dr Benson Ogunjimi.

"Our findings should allay some fears about implementing childhood chickenpox vaccination," he says.

Vaccination programmes are established in some countries including the US, Australia, Greece, Germany, Japan and Taiwan. However, as most - but not all - chickenpox cases are mild, whereas [shingles](#) causes a painful rash and can lead to complications such as prolonged pain (postherpetic neuralgia), policy makers in other countries have hesitated to introduce childhood vaccination.

Shingles occurs most often in individuals with a declined immunological status, such as HIV or cancer patients. Ageing is also assumed to increase susceptibility. In previous studies, Ogunjimi and colleagues found that the impact of ageing is exacerbated by Cytomegalovirus infection, another virus in the herpes family.

The current model combines data from individuals relating to all these factors, making it more accurate than previous models and easier to verify with future clinical studies.

"Our [new model](#) relates to real biological functions, increasing its value for determining health policy," says Dr Ogunjimi.

Provided by eLife

Citation: Chickenpox vaccination does increase shingles cases, but mainly in young adults (2015, August 11) retrieved 6 May 2024 from <https://medicalxpress.com/news/2015-08-chickenpox->

[vaccination-shingles-cases-young.html](https://www.medicalxpress.com/vaccination-shingles-cases-young.html)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.