

# Depressed teens at risk of heart disease, early monitoring urged

August 10 2015

---



Credit: George Hodan/Public Domain

For the first time, experts urge early monitoring for heart and blood vessel disease among teens with major depression or bipolar disorder, according to an American Heart Association scientific statement.

"Major Depressive Disorder and Bipolar Disorder Predispose Youth to Accelerated Atherosclerosis and Early Cardiovascular Disease" is

published in *Circulation*, a journal of the American Heart Association.

This statement makes recommendations to consider these mood disorders as independent, moderate [risk factors](#) for cardiovascular diseases and is based on a group of recent scientific studies including those that reported cardiovascular events such as heart attacks and deaths among young people. For example, a 2011 population study of more than 7,000 U.S. young adults under age 30 found that a history of depression or an attempted suicide was the number one risk factor for [heart disease](#) death caused by narrowed/clogged arteries in young women, and the number 4 risk factor in young men.

"Youth with mood disorders are not yet widely recognized as a group at increased risk for excessive and early heart disease. We hope these guidelines will spur action from patients, families and healthcare providers to reduce the risk of cardiovascular disease among these youth," said Benjamin I. Goldstein, M.D., Ph.D., lead author of the statement and a child-adolescent psychiatrist at Sunnybrook Health Sciences Centre and the University of Toronto, in Toronto, Ontario, Canada.

Major depression and [bipolar disorder](#) are common mood disorders, affecting about 10 percent of U.S. adolescents. Worldwide, [major depression](#) is the main cause of disability among teens, while bipolar disorder is the fourth-leading cause. Symptoms of major depression include persistent feelings of sadness and loss of interest in previously enjoyed activities. Bipolar disorder can cause severe mood swings between periods of major depression and mania, which may include increased elation, energy, irritability and a decreased need for sleep.

Previous research showed that adults with major depression and bipolar disorder are far more likely to have cardiovascular disease, and that they experience cardiovascular disease at much earlier ages than other adults.

Since cardiovascular disease may begin early in life, the authors want to increase awareness and recognition of mood disorders among young people as moderate-risk conditions for early cardiovascular disease.

After systematically analyzing published research, the authors found that teens with major depression or bipolar disorder are more likely than other teens to have several [cardiovascular disease risk](#) factors including:

- [high blood pressure](#);
- high cholesterol;
- obesity, especially around the midsection;
- type 2 diabetes;
- and hardening of the arteries.

The biological causes of these increased risks remain unclear, but they may be related to inflammation and other types of cell damage, which some studies found occur more frequently among teens with mood disorders compared to other teens.

While teens with [mood disorders](#) were more likely than other teens to have unhealthy behaviors, like drug abuse, smoking and being physically inactive, those factors alone do not explain the increased [cardiovascular risk](#).

Similarly, medications do not fully explain the increased risk. While certain mood medications can cause weight gain, and high blood pressure, cholesterol and blood-sugar levels, most of the teens in the studies analyzed were not medicated.

Based on these findings, the authors advise including major depression and bipolar disorder as a moderate risk factor for [cardiovascular disease](#) in teens.

"Mood disorders are often lifelong conditions, and managing cardiovascular risk early and assertively is tremendously important if we are to be successful in ensuring that the next generation of youth has better cardiovascular outcomes," Goldstein said. "These disorders indicate an increased risk of heart disease that requires increased vigilance and action at the earliest possible stage."

Provided by American Heart Association

Citation: Depressed teens at risk of heart disease, early monitoring urged (2015, August 10) retrieved 6 May 2024 from

<https://medicalxpress.com/news/2015-08-depressed-teens-heart-disease-early.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--