

# New diagnostic tools for dehydration severity in children

August 21 2015, by David Orenstein

---



Nurse Ilishabeth Samadder sees many young children at the International Centre for Diarrhoeal Disease Research in Bangladesh. Credit: Dr. Adam Levine

Dehydration from diarrhea, either from viral infection or cholera,

accounts for 10 percent of all child deaths worldwide. Though it's a pervasive problem, there are no clinical guidelines that perform better than chance for assessing dehydration severity. In a new study, researchers report that they statistically derived two simple diagnostic tools with high sensitivity and specificity from the cases of 771 children aged 0 to 5 in Dhaka, Bangladesh.

Named for the city where thousands of children suffer diarrhea-related [dehydration](#) every spring as cholera cases mount, the Dehydration: Assessing Kids Accurately (DHAKA) system includes both a simple scoring tool and a decision tree that clinicians can quickly apply when young patients arrive at a clinic, said study lead author Dr. Adam Levine, assistant professor of emergency medicine in the Alpert Medical School of Brown University and a physician at Rhode Island Hospital. Either tool requires observing only a few symptoms that are readily apparent in a young child.

Assessing dehydration severity is crucial in resource-constrained clinics across the developing world. Clearly, undertreating a severe case has a cost, but so does overtreating a milder one. Clinics need to reserve intravenous fluids for the most severely affected children. Less sick children can be rehydrated orally.

"You could miss kids with severe dehydration and not apply the correct treatment, and you could overtreat kids who don't need it, which wastes precious resources," Levine said. "Diagnosis matters not only for the health of the individual child but also from a public health and health economics standpoint."

In past studies, Levine and other researchers evaluated the performance of existing diagnostic systems, including the most prominent one developed by the World Health Organization. Levine analyzed the cases of 178 children in Rwanda [[LINK](#)]. He and others found that existing

systems statistically didn't perform better than chance.

## Children too sick for tears


Clinical Sign	Finding	Points
General appearance	Normal	0
	Restless/irritable	2
	Lethargic/ unconscious	4
Respirations	Normal	0
	Deep	2
Skin pinch	Normal	0
	Slow	2
	Very slow	4
Tears	Normal	0
	Decreased	1
	Absent	2
DHAKA Dehydration Score Categories		Points
No dehydration		0–1
Some dehydration		2–3
Severe dehydration		≥4

By assessing a few symptoms, a doctor or nurse could add up a simple score to arrive at a diagnosis. Credit: Dr. Adam Levine

Levine wanted to develop something that could. The mission brought him to the International Centre for Diarrhoeal Disease Research in Bangladesh in the spring of 2014. The center sees so many cases that Levine and his co-authors could gather a lot of data on the symptoms children exhibit when they first present at the center and the eventual confirmed degree of their dehydration (as measured by how much fluid they needed to return to full health).

In the study, Levine and his colleagues tracked 10 potential symptoms to see which ones were accurately predictive of true dehydration severity. The analysis found that just a few were needed to produce the combination of [high sensitivity](#) (does it detect dehydration severity?) and high specificity (does it rule out other possible problems?) needed to provide an accurate diagnosis.

For the DHAKA Dehydration Score these symptoms are the general appearance and demeanor of a child, how quickly skin returns to smooth after being pinched, whether breathing is unusually deep, and whether tears are absent or only barely present when the child is crying. A severely dehydrated child has no water to spare for tears.



To arrive at a diagnosis, clinicians can opt to follow a two-step decision tree. Credit: Dr. Adam Levine

For the DHAKA Dehydration Decision Tree, a two-step flowchart process, the symptoms were general appearance and, depending on that, either the skin pinch or whether the child's eyes appeared sunken.

When analyzed to see how predictive they were of dehydration severity, the tools scored well, Levine said. On an overall measure of accuracy called AUC, which ranges from 0 to 1, the score rated a 0.8 and the tree rated a 0.76 for severe dehydration. The score's sensitivity, on a percentage scale, was 83 percent for severe dehydration and its specificity was 57 percent. For the tree, sensitivity was 81 percent and specificity was 67 percent.

The next step was to further validate the tools by testing them in clinical practice with a new population of children. That occurred this spring at

the center, and analysis of the results is pending.

"The goal is that hopefully the DHAKA score or the DHAKA tree will validate well in this new population," Levine said. "If it does, then we would say that it should be used clinically going forward as the standard of care."

Provided by Brown University

Citation: New diagnostic tools for dehydration severity in children (2015, August 21) retrieved 4 May 2024 from <https://medicalxpress.com/news/2015-08-diagnostic-tools-dehydration-severity-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.