


Engineered hot fat implants reduce weight gain in mice

August 20 2015



A thermograph of brown-like fat implanted in anesthetized animal at room temperature. Implant, in area A, is significantly warmer (30.86 degrees Celsius maximum) compared with control region in area B (29.43 degrees Celsius maximum). Credit: Kevin Tharp and Andreas Stahl

Scientists at the University of California, Berkeley, have developed a novel way to engineer the growth and expansion of energy-burning "good" fat, and then found that this fat helped reduce weight gain and

lower blood glucose levels in mice.

The authors of the study, to be published Thursday, Aug. 20, in the journal *Diabetes*, said their technique could eventually lead to new approaches to combat obesity, diabetes and other metabolic disorders.

The researchers used a specifically tailored hydrogel to "scaffold" and control an implant containing [stem cells](#) to form a functional [brown-fat](#)-like tissue. While white fat—the kind associated with obesity—stores excess energy, brown fat serves as a heat generator, burning calories as it does its job.


"What is truly exciting about this system is its potential to provide plentiful supplies of brown fat for therapeutic purposes," said study lead author Kevin Tharp, a Ph.D. student in the Department of Nutritional Sciences and Toxicology. "The implant is made from the stem cells that reside in white fat, which could be made from tissue obtained through liposuction."

Human babies, which cannot yet produce heat by shivering, have greater stores of brown fat, so-called because it contains high levels of darker-hued mitochondria. It was once believed that brown fat disappears with age, but in recent years, this tissue has been discovered in the neck, shoulders, and spinal cord among adults.

"This is figuratively and literally a hot area of research right now," said the study's senior author, Andreas Stahl, an associate professor of nutritional sciences and toxicology. "We are the first to implant in mice an artificial brown-fat depot and show that it has the expected effects on body temperature and beneficial effects on metabolism."

Studies have shown that cold temperatures can bump up activity in brown fat. Stahl noted, however, that the exposure to cold often led to

increases in food intake, as well, potentially negating any calorie-burning benefits from brown-fat activity.



The appearance of an implant, less than a centimeter long, removed after two weeks. Notably, the fat shows the development of blood vessels and a marked brown hue. Credit: Kevin Tharp and Andreas Stahl


Get the fat, not the cold

There are three basic types of fat tissue in our bodies. There is the classic, energy-storing white fat that many of us are most familiar with, and two kinds of energy-burning fat used to generate heat, namely brown fat, which arises during fetal development, and [beige fat](#), which is brown-like fat formed within white fat tissue after exposure to cold and other situations.

This UC Berkeley experiment explored the idea of increasing brown-like beige fat without the temperature drop. Stahl teamed up with Kevin Healy, UC Berkeley professor of bioengineering, and postdoctoral researcher Amit Jha to develop a system of physical cues to guide stem cell differentiation.

"It's already known that for a number of organs, including the heart, the extracellular matrix in which a cell resides provides signals to guide growth and development," said Healy, who also has an appointment in the Department of Materials Science and Engineering. "We applied this concept to stem cells isolated from white-fat tissue."

The specific matrix recipe for converting white-fat stem cells to brown fat had been unclear, the researchers said, but they noted that previous studies suggested that stiffness of the surrounding environment was a factor. White-fat stem cells placed in a 3D environment that is soft, with little resistance as the cell grows, became fat. When the surrounding environment was very stiff, the stem cells grew into bone.



Implants respond to a stimulated cold challenge with increased fat uptake. Images on top are anesthetized animals with implants that emit light proportionally to the uptake rate of fatty acids. The bar graphs show the higher rate of activity under stimulated conditions. Credit: Kevin Tharp and Andreas Stahl

Injected brown fat reduces weight gain, glucose

The researchers created a tightly knit 3D mesh in a hydrogel containing water, hyaluronic acid and short protein sequences associated with brown-fat growth and function. Hyaluronic acid is a naturally occurring carbohydrate that helps make water thicker and gel-like. They then took white-fat stem cells from mice genetically engineered to express an enzyme from fireflies. This made the cells luminescent, allowing the researchers to track them more easily.

The researchers then added the cells to the hydrogel and, before the mixture thickened, injected them under the skin of genetically identical mice.

The gel polymerizes after injection and completes its transformation in the animal. The researchers monitored the glowing cells after injection to determine how well they stayed put, how long they persisted in the body and whether they were metabolically functional.

They noticed an increase in the core body temperature of the mice at ambient temperatures of 21 degrees Celsius and after 24 hours at a chilly 4 degrees Celsius. In both cases, the mice with the implanted cells were up to half a degree Celsius warmer than a control group of mice with no injection. The higher the concentration of cells, the larger the effect on temperature.

The researchers also put the experimental mice on a high-fat diet. By the end of three weeks, the mice with injected beige fat gained half as much weight and had lower levels of blood glucose and circulating fatty acids compared with control mice.

"This is a feasibility study, but the results were very encouraging," said Stahl. "It is the first time an optimized 3D environment has been created to stimulate the growth of brown-like fat. Given the negative health effects of obesity, research into the role of brown fat should continue to see if these findings would be effective in humans."

Provided by University of California - Berkeley

Citation: Engineered hot fat implants reduce weight gain in mice (2015, August 20) retrieved 4 May 2024 from <https://medicalxpress.com/news/2015-08-hot-fat-implants-weight-gain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.