

Long-term brain changes persist years after drug abuse and recovery

August 18 2015



It's known that brain changes are present in drug addicts even when they have been abstinent for a short period of time. Now new research shows that alterations persist in long-term abstinent heroin-dependent individuals as well.

Through the use of functional magnetic resonance imaging, investigators analyzed the brains of 30 heroin-addicted individuals after a long period of abstinence (more than 3 years) and compared the results with those of 30 healthy controls.

The team found that in the former [heroin users](#), there was significant dysfunctional activity in the nucleus accumbens region of the brain, an

area that is involved in appetitive drive and reward-seeking behavior.

The *Journal of Neuroscience Research* findings suggests that intense craving for drugs may be triggered in the presence of stressors or other environmental cues in individuals who have successfully remained abstinent for long periods of time. Taking this into consideration may aid the development of treatment and rehabilitation strategies for those suffering from drug addiction.

More information: "Abnormal resting-state functional connectivity of the nucleus accumbens in multi-year abstinent heroin addicts." [DOI: 10.1002/jnr.23608](https://doi.org/10.1002/jnr.23608)

Provided by Wiley

Citation: Long-term brain changes persist years after drug abuse and recovery (2015, August 18) retrieved 27 April 2024 from <https://medicalxpress.com/news/2015-08-long-term-brain-persist-years-drug.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.