


miR-7 suppresses stomach cancer

August 10 2015



Zhao et al. identify a regulatory circuit in human gastric cells that becomes compromised by *H. pylori*-induced inflammation and leads to the development of stomach cancer. Credit: Zhao et al., 2015

A study in *The Journal of Cell Biology* reveals that the microRNA miR-7 suppresses gastric (stomach) cancer by inhibiting a key signaling pathway, and that this protective mechanism is compromised by the cancer-causing bacterium *H. pylori*. Finding drugs capable of inducing miR-7 could therefore prove to be an effective treatment against the progression of gastric cancer.

Gastric cancer is the fourth most common cancer and the third leading cause of cancer-related deaths worldwide, according to the National Institutes of Health. miR-7, which is frequently decreased in gastric cancers, can stop the [cancer cells](#) from spreading to other tissues by inhibiting a particular growth factor receptor (called IGF1R). Whether miR-7 also suppresses earlier stages of gastric cancer is unknown, however, so researchers in China screened for new targets of the microRNA.

Dai-Ming Fan and colleagues found that miR-7 directly targets the genes *RELA* and *FOS*, which encode proteins involved in the pro-oncogenic NF- κ B and AP-1 signaling pathways, respectively. In human gastric cancer samples, low miR-7 levels correlated with elevated levels of *RELA* and *FOS* proteins and poor patient survival. Increasing levels of miR-7 reduced *RELA* and *FOS* levels and inhibited tumor growth in mice.

The researchers found that, as well as directly suppressing *RELA* expression, miR-7 could control the protein's activation by targeting its upstream kinase (IKK ϵ) in the NF- κ B pathway. Yet, this same pathway was itself able to repress miR-7 expression, indicating that miR-7 would be unable to restrain *RELA*'s activity if the NF- κ B pathway were strongly activated.

Chronic *H. pylori* infection is a major risk factor for [gastric cancer](#), in part because the bacterium can hyperactivate the NF- κ B pathway.

Accordingly, Fan and colleagues found that culturing *H. pylori* with gastric cells activated IKK ϵ and RELA, and reduced the expression of miR-7, a potentially key step in the transformation of healthy gastric cells into malignant ones.

More information: Zhao, X.-D., et al. 2015. *J. Cell Biol.*
[dx.doi.org/10.1083/jcb.201501073](https://doi.org/10.1083/jcb.201501073)

Provided by Rockefeller University Press

Citation: miR-7 suppresses stomach cancer (2015, August 10) retrieved 25 April 2024 from
<https://medicalxpress.com/news/2015-08-mir-suppresses-stomach-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.