

Understanding the molecular mechanism leading to addiction in humans

August 4 2015

- A new study just published in the prestigious journal *Molecular Psychiatry* by the team led by Salah El Mestikawy, Ph.D., researcher at the Douglas Mental Health University Institute (CIUSSS de l'Ouest-de-l'île-de-Montréal), professor at McGill University and head of research at CNRS INSERM UPMC in Paris, opens the field to new understanding of the molecular mechanism underlying addiction in humans.

This work was accomplished through collaboration with several international research teams. Among these researchers was Stéphane Jamain (CR1 INSERM, Université de Créteil, France), a French geneticist specialized in psychiatric disorders.

The research explained in simple terms

To communicate with each other, brain cells ([neurons](#)) use chemicals known as neurotransmitters. Among the classical neurotransmitters are dopamine, serotonin, acetylcholine and glutamate. Until recently, it was believed that a given neuron used only one classical [neurotransmitter](#). Some time ago, Salah El Mestikawy's team contributed to the surprising discovery that subpopulations of neurons are able to use two transmitters. In other words, unlike the vast majority of neurons, these neurons are bilingual. Most neurons in the brain respond with either "yes", "no" or "maybe". El Mestikawy and his colleagues found that in order to regulate our reward behaviours, a small population of neurons in our reward centre are able to respond with both "yes" and "no". These

neurons are known as cholinergic neurons, while the reward centre is known as the nucleus accumbens. In this new work, the researchers discovered that when they shut down one of the languages (the "no") of [cholinergic neurons](#) within the reward centre, mice showed a marked predilection for cocaine. This work explains the [molecular mechanism](#) underlying reward behaviour, while identifying an unsuspected target in the treatment of addiction.

Finally and perhaps even more important, the team found that humans with mutation of a key gene for this co-transmission were 10 times more vulnerable to very severe forms of addiction.

"These discoveries are bringing about a major transformation in the field of addiction. We are beginning to decipher and understand the complex regulation of basic behaviours" says Salah El Mestikawy.

Provided by Douglas Mental Health University Institute

Citation: Understanding the molecular mechanism leading to addiction in humans (2015, August 4) retrieved 5 May 2024 from <https://medicalxpress.com/news/2015-08-molecular-mechanism-addiction-humans.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--