

Physician support key to successful weight loss, study shows

August 21 2015

A review of survey data from more than 300 obese people who participated in a federally funded weight loss clinical trial found that although the overall weight loss rates were modest, those who rated their primary care doctor's support as particularly helpful lost about twice as many pounds as those who didn't.

In a report on the study by Johns Hopkins researchers, published in the Aug. 21 issue of *Patient Education and Counseling*, the researchers say the findings could inform the development of weight loss programs that give [primary care](#) physicians a starring role.

Researchers have long known that high-quality patient-doctor relationships marked by empathy, good communication, collaboration and trust are linked to better adherence to medication schedules, appointment keeping and other good outcomes, says Wendy L. Bennett, M.D., M.P.H., assistant professor of medicine at the Johns Hopkins University School of Medicine and a [primary care physician](#) at Johns Hopkins Bayview Medical Center. Previous studies also have shown, she says, that obese patients are more likely to report poor physician-patient relationships, with evidence of decreased respect and weight bias from providers.

To see whether and what aspects of those relationships might influence weight loss efforts, Bennett and her colleagues reviewed information gathered by Johns Hopkins' Practice-based Opportunities for Weight Reduction ([POWER](#)) trial, a two-year, randomized, controlled study

funded by the federal government. During the trial, some [obese patients](#) worked to lose weight with the aid of health coaches while their efforts were supervised by their primary care physicians.

At the end of the trial, patients filled out surveys that asked, in part, about their relationships with their primary care physician, including questions about how often their providers explained things clearly, listened carefully and showed respect, as well as how helpful their physicians' involvement was in the trial. Of the 347 patients who filled out surveys, about 63 percent were female, about 40 percent were African-American and all were obese, with body mass indices of 36.3 on average. Each participant also had one of three cardiovascular disease risk factors: high blood pressure, high cholesterol or diabetes.

Results of a review showed that nearly all of the 347 patient surveys reviewed for the Johns Hopkins study reported high-quality relationships with their physicians, with the overall relationship showing little effect on weight loss. However, those patients who gave their physicians the highest ratings on "helpfulness" during the trial lost an average of 11 pounds, compared to just over 5 pounds for those who gave their physicians the lowest "helpfulness" ratings.

Current National Institutes of Health statistics suggest that more than one-third of adults in the United States are obese. Though Medicare and private insurance reimbursements are low or nonexistent for physician-guided weight loss interventions, Bennett says, the findings could spur new reimbursement models that provide for physician involvement and enable more team-based care models.

"This trial supports other evidence that providers are very important in their patients' [weight loss efforts](#)," Bennett says. Many current weight loss programs are commercially run, she adds, and patients often join these programs without their physician's knowledge.

"Incorporating physicians into future programs might lead [patients](#) to more successful [weight loss](#)," she says.

Provided by Johns Hopkins University School of Medicine

Citation: Physician support key to successful weight loss, study shows (2015, August 21)
retrieved 25 April 2024 from

<https://medicalxpress.com/news/2015-08-physician-key-successful-weight-loss.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.