

Speedy treatment with new clot retrieval device vastly improves stroke outcomes

August 18 2015, by Kim Irwin



Micrograph showing cortical pseudolaminar necrosis, a finding seen in strokes on medical imaging and at autopsy. H&E-LFB stain. Credit: Nephron/Wikipedia

In the treatment of stroke patients, time really is brain. A few minutes can mean the difference between patients living independently or suffering debilitating disabilities.

Now, UCLA researchers have shown, for the first time, that speedy



treatment with a new-generation stent clot retrieval device results in greatly improved outcomes, and that even a five-minute delay negatively affects patients.

The one-year study found that when <u>blood flow</u> was restored to the brain within four hours of the start of a stroke, 80 percent of patients had a very good outcome—meaning that they survived and were able to live independently three months later.

The researchers also found that the odds of a poorer outcome increased by one percent for every five minutes that passed between the onset of a stroke and the time when doctors reopened the blocked artery, according to Dr. Sunil Sheth, a fellow in radiological sciences at the David Geffen School of Medicine at UCLA.

"The impact of the speed of treatment was found to be enormous," Sheth said. "These findings indicate that <u>stroke centers</u> worldwide should strive to bring patients quickly to comprehensive stroke centers offering this therapy, and that those centers should work to shorten the times from patient arrival to artery reopening."

The study results appear today in the peer-reviewed journal *Annals of Neurology*.

In recent years, doctors' ability to restore blood flow to a patient's brain after a stroke has advanced tremendously, thanks to the development of stent retrieval devices that pull clots out of the blocked artery.

Each device consists of a thin mesh column that is introduced into a blood vessel in the leg and guided up to the blocked vessel in the brain, where it captures and removes the clot to restore blood flow to the brain.

The new-generation devices have an improved design that allows them to



more easily surround and remove the clot. The first clot retrieval device for stroke was invented at UCLA.

"When it comes to treating stroke, every minute counts," said study senior author Dr. Jeffrey Saver, director of the UCLA Stroke Center. "The faster you can restore the blood flow to normal, the better the chances of a good recovery. Our study shows that the currently used stent clot retrieval devices are a tremendous improvement from the older technologies."

Saver said a number of changes should be implemented immediately across the entire spectrum of stroke care—including improving public awareness about the signs and symptoms of stroke, educating emergency medical crews to route patients to the appropriate medical centers, and monitoring the quality of care that people receive in the hospital.

Other studies have looked at the impact of speed of clot retrieval, but those studies analyzed data from outdated trials that used obsolete techniques and devices. The new research is the first to use data from the version of the devices currently in use, Sheth said.

In addition, Sheth said, the UCLA team took a more refined look at how <u>stroke patients</u> fared after receiving treatment. Instead of lumping them into two groups—"good" and "bad"—as previous studies did, the team used the results of detailed patient interviews and neurological exams to divide the patients into seven different categories, which more accurately reflect people's experiences of recovering from stroke.

Even when doctors are able to clear the blocked blood vessel quickly, not all patients recover well. In future research, the UCLA team will focus on ways to identify which patients are at an increased risk of poor outcomes, in order to improve doctors' ability to provide accurate prognostic information to them and their families.



According to the American Heart Association and the American Stroke Association, stroke is the fourth most common cause of death and a leading cause of adult disability in the United States. On average, one American suffers a stroke every 40 seconds and one dies of a stroke every four minutes. It is estimated that 795,000 Americans suffer a new or recurrent <u>stroke</u> each year.

"As clinicians we are constantly reminded of the challenges brought on by neurological deficits in our <u>patients</u>," the study says. "More granular analyses such as these that are sensitive to detect smaller changes in disability allow for richer discussions when advising on treatment approaches, as well as providing counseling in the acute setting."

More information: "Time to endovascular reperfusion and degree of disability in acute stroke." *Ann Neurol.*. doi: 10.1002/ana.24474

Provided by University of California, Los Angeles

Citation: Speedy treatment with new clot retrieval device vastly improves stroke outcomes (2015, August 18) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2015-08-speedy-treatment-clot-device-vastly.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.