

Young adults living in polluted city show early signs of cardiovascular risk

August 30 2015

Young adults living in a polluted city show early signs of cardiovascular risk, according to research presented at ESC Congress today by Dr Krzysztof Bryniarski from Jagiellonian University, Collegium Medicum in Krakow, Poland. Residing in a polluted city was associated with higher levels of inflammatory markers in otherwise healthy adolescents and young adults, which indicate a greater risk of having a heart attack in future.

Dr Bryniarski said: "This study was conducted by a group of medical students from Jagiellonian University in Krakow, supervised by Professors T. Guzik and A. Wysokinski, who wanted to know what impact long term air pollution is having on the cardiovascular health of [young adults](#). Krakow is one of the most polluted cities in Europe. We decided to compare some markers of cardiovascular risk in our city with risk in Lublin, which has similar general demographic characteristics but much lower traffic congestion and two fold lower air pollution."

The current study assessed the association between long term air pollution exposure and blood pressure and [inflammatory markers](#) in Krakow and Lublin, two large Polish cities each with more than 300 000 inhabitants. It included 826 randomly selected healthy young adults aged 16 to 22 years (average age 18), of whom 444 lived in Krakow and 382 lived in Lublin. Participants came from similar types of schools and social backgrounds and had lived in their home city since birth.

The researchers measured blood pressure, heart rate and blood levels of

inflammatory markers including C-reactive protein (CRP), high sensitivity CRP (hsCRP), homocysteine and fibrinogen which are early indicators of cardiovascular disease. They also assessed numerous potential confounding factors such as body mass index (BMI), lifestyle, ethnicity and family history.

Data on air pollution was obtained from air quality monitoring stations. Average 10 year air pollution levels were nearly double in Krakow compared to Lublin. Average PM_{2.5} levels for example were 41.7 µg/m³ in Krakow and 22.4 µg/m³ in Lublin while PM₁₀ was 56.9 µg/m³ in Krakow and 29.4 µg/m³ in Lublin.

The investigators found significantly higher levels of CRP, hsCRP, homocysteine and fibrinogen levels in Krakow residents compared to Lublin residents (p

Citation: Young adults living in polluted city show early signs of cardiovascular risk (2015, August 30) retrieved 17 April 2024 from <https://medicalxpress.com/news/2015-08-young-adults-polluted-city-early.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--