

Excessive daytime sleepiness and long naps linked to increased diabetes risk

September 17 2015

New research presented at this year's annual meeting of the European Association for the Study of Diabetes (EASD) shows that daytime sleepiness and taking long naps during the day are both associated with an increased risk of type 2 diabetes. The research is by Dr Tomohide Yamada, University of Tokyo, Japan.

Sleep is an important component of a healthy life, along with a good diet and appropriate physical activity. Excessive daytime sleepiness is widely prevalent around the world, as is the habit of taking short sleeps or "napping". Daytime naps are usually brief, but can range from a few minutes to a few hours. The frequency varies from taking an occasional nap to planned rest periods several times daily for habitual nappers. Some individuals take a nap because they are excessively sleepy during the daytime as a result of a sleep disorder.

In this new study, the authors did a meta-analysis to investigate the association between daytime sleepiness or napping and the risk of type 2 [diabetes](#). They searched Medline, the Cochrane Library, and Web of Science for articles published up to November 2014 using the keywords daytime sleepiness, nap, and diabetes. Among 683 studies initially identified, a total of 10 were deemed of good quality and included 261,365 Asian and Western subjects. The studies came from Sweden, Spain, Finland, and Germany (daytime sleepiness) and the USA, China and Germany (napping). Excessive daytime sleepiness was defined as answering yes to questions like "Do you have a problem with sleepiness during the daytime?". Daytime napping was defined on the basis of

answering yes to questions such as "Do you take a daytime nap?" or "Do you sleep during the day?".

Excessive daytime sleepiness was found to increase the risk of diabetes by 56%, while a longer daytime nap of 60 minutes or more increased the risk by 46%. In contrast, a shorter nap (60 mins or less per day) did not increase the risk of diabetes. The analysis showed there was no effect of napping up to about 40 minutes per day, after which risk began to increase sharply.

The authors conclude: "Excessive [daytime sleepiness](#) and taking longer naps were associated with increased risk of type 2 diabetes, with a short nap not increasing this risk."

They add: "Daytime napping might be a consequence of night-time sleep disturbance such as obstructive sleep apnoea (OSA). Epidemiological studies have shown that obstructive sleep apnoea is independently linked to blockages (ischaemia) of heart arteries, stroke, fatal and non-fatal cardiovascular events, and all-cause mortality. "

They explain further: "Several studies have demonstrated the beneficial effects of taking short naps less than 30 minutes in duration, which help to increase alertness and motor skills. A short nap finishes before the onset of deep slow-wave sleep. Entering deep slow-wave sleep and then failing to complete the normal sleep cycle can result in a phenomenon known as sleep inertia, in which a person feels groggy, disoriented, and even sleepier than before napping. Although the mechanisms by which a short [nap](#) might decrease the risk of diabetes are still unclear, such duration-dependent differences in the effects of sleep might partly explain our findings."

Provided by Diabetologia

Citation: Excessive daytime sleepiness and long naps linked to increased diabetes risk (2015, September 17) retrieved 5 May 2024 from <https://medicalxpress.com/news/2015-09-excessive-daytime-sleepiness-naps-linked.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.