

Hematologic markers predict nasopharyngeal cancer survival

September 22 2015



(HealthDay)—Pretreatment neutrophil-lymphocyte ratio (NLR) and platelet-lymphocyte ratio (PLR) may be predictors for survival in nasopharyngeal carcinoma, according to a study published online Sept. 11 in *Head & Neck*.

Wei Sun, M.D., from the Huazhong University of Science and Technology in Wuhan, China, and colleagues evaluated the prognostic significance of various hematologic parameters in 251 patients with [nasopharyngeal carcinoma](#).

The researchers found that $NLR \geq 2.7$ (hazard ratio, 2.01; $P = 0.005$) and $PLR \geq 167.2$ (hazard ratio, 2.12; $P = 0.001$) were significantly associated with shorter progression-free survival, while $PLR \geq 163.4$ was correlated

with poor overall survival (hazard ratio, 2.64; P = 0.011).

"Pretreatment NLR and PLR can be independent prognostic factors for patients with NPC," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Hematologic markers predict nasopharyngeal cancer survival (2015, September 22)
retrieved 24 April 2024 from
<https://medicalxpress.com/news/2015-09-hematologic-markers-nasopharyngeal-cancer-survival.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--