

Incidence of diabetes and pre-diabetes on the rise

September 17 2015, by Scott Gilbert



Credit: AI-generated image ([disclaimer](#))

A generation ago, there was no such thing as pre-diabetes. Well, the condition existed – it just wasn't given a name or diagnosed the way it is now.

Dr. Chris Fan, practice site director for the endocrinology and

nephrology clinic at Penn State Hershey Medical Center, said news reports that nearly half of all Americans are either diabetic or pre-diabetic are pretty accurate.

"If you figure that 7 to 10 percent of Americans have diabetes, another 7 to 10 percent have it and don't realize it, and the number of people who have pre-diabetes is at least double that, you're not far from half," Fan said.

Although the definition of diabetes and the criteria for being diagnosed have not changed, more Americans are living – or will soon live with – the chronic condition.

"As our collective waistline expands, more people are becoming diabetic or pre-diabetic," Fan said. Another challenge is that not everyone is diligent about going to their family doctor for screenings and blood tests that can indicate the disease. "A lot of times people don't even realize they have the condition," he said.

As with most health issues, early diagnosis and treatment can prevent complications. "It's natural for people not to want to take drugs – especially if it involves an injection," Fan said. "But you're better off starting treatment early than waiting until you have no choice."

He points out that Type 2 diabetes – the type that affects 95 percent of those with diabetes – is associated with a number of complications that shorten life expectancy or make life "considerably less pleasant."

Fan said diabetes is the leading cause of nontraumatic limb amputations and is the principal reason people go on dialysis and need kidney transplants. It is the leading cause of blindness in adults under 75. Being diabetic also increases a person's risk for heart attack and stroke "by at least twofold, and possibly as much as tenfold."

Diabetes is defined as a fasting blood sugar of more than 125, a random blood sugar of 200 or more with classic symptoms, a three-month mean sugar (also known as hemoglobin A1C) of 6.5 percent or more, or a blood sugar of 200 or more after a two-hour glucose tolerance test.

A fasting [blood sugar](#) of 100 to 125, a two-hour glucose tolerance test result of 140 to 199, or an A1C between 5.7 and 6.4 percent denotes pre-diabetes. By comparison, fasting blood sugars in the 70s to 90s are considered to be normal.

Fan said time spent with a diabetic dietician and nurse educator are important for pre-diabetics looking to prevent progression of their condition. "Unfortunately, there is a relative shortage of those individuals and many insurances generally won't cover a visit to a dietician because the system isn't set up to prevent disease, but to treat it once it occurs," Fan said.

His hope is that a diagnosis of pre-[diabetes](#) will galvanize people to action, whether that means incorporating exercise and/or healthier food choices into their daily routines to lose weight and decrease their risk. "It's important to know if you have it, if for no other reason than it allows you to take steps to revert to normal or prevent progressing."

Provided by Pennsylvania State University

Citation: Incidence of diabetes and pre-diabetes on the rise (2015, September 17) retrieved 10 April 2024 from <https://medicalxpress.com/news/2015-09-incidence-diabetes-pre-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
