


Preoperative breast MRI use increased from 2003 to 2012

September 25 2015



(HealthDay)—From 2003 to 2012 there was a significant increase in preoperative breast magnetic resonance imaging (MRI) use in women with breast cancer, according to a study published online Sept. 24 in *JAMA Oncology*.

Angel Arnaout, M.D., from Ottawa Hospital in Canada, and colleagues describe patterns of preoperative breast MRI utilization in a [health care](#) system with universal insurance. Data were included from administrative health care databases in Ontario, Canada, in a [retrospective cohort study](#). The cohort included 53,015 patients with primary operable [breast cancer](#) who were treated from 2003 to 2012.

The researchers found that 14.8 percent of patients underwent a preoperative MRI. There was an eight-fold increase in MRI use across all stages during the 10-year study period (from 3 to 24 percent; P

"Preoperative breast MRI use has increased substantially in routine clinical practice and is associated with a significant increase in ancillary investigations, wait time to surgery, mastectomies, and contralateral prophylactic mastectomies," the authors write.

More information: [Abstract](#)

[Full Text](#)

[Editorial](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Preoperative breast MRI use increased from 2003 to 2012 (2015, September 25)
retrieved 18 April 2024 from

<https://medicalxpress.com/news/2015-09-preoperative-breast-mri.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.