

# Substantial variability among clinicians in imaging recs

September 17 2015

---



(HealthDay)—There is substantial variability between radiologists and technologists in recommendation rates for additional imaging (RAI) during ultrasound interpretation, according to a study published in the October issue of the *Journal of Clinical Ultrasound*.

Nathaniel E. Margolis, M.D., from New York University Langone Medical Center, and colleagues retrospectively reviewed 719 [ultrasound](#) reports from a single academic medical center for the presence of RAI. One of three abdominal radiologists had interpreted all studies. Examinations were performed at either an outpatient radiology facility with no on-site radiologist (n=299) or at an inpatient emergency department or hospital-based outpatient setting that had an on-site radiologist (n=420).

The researchers found that there were significant differences between each pair of radiologists in terms of overall frequency of RAI (P radiologist location (P = 0.15-0.93).

"The observed substantial variability in RAI between radiologists and technologists warrants further study, with consideration of strategies to optimize RAI within ultrasound reports," the authors write.

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Substantial variability among clinicians in imaging recs (2015, September 17) retrieved 30 April 2024 from  
<https://medicalxpress.com/news/2015-09-substantial-variability-clinicians-imaging-recs.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--