

# In terminally ill patients, some types of delirium are a sign of 'imminent death'

September 23 2015

---

In cancer patients nearing the end of life, certain subtypes of delirium—specifically, hypoactive and "mixed" delirium—are a strong indicator that death will come soon, reports a study in *Psychosomatic Medicine: Journal of Biobehavioral Medicine*, the official journal of the American Psychosomatic Society.

"Terminally ill [patients](#) with the hypoactive or mixed subtypes of delirium showed a higher probability of imminent death, with even earlier mortality among younger patients," according to the new research by Sung-Wan Kim, MD, and colleagues of Chonnam National University Medical School Gwangju, Republic of Korea. They believe their findings might help make more accurate predictions of survival in patients nearing the end of life.

## Shorter Survival in Patients with Hypoactive/Mixed Delirium

The researchers looked at the relationship between delirium and survival time in 322 patients with terminal cancer entering [palliative care](#). Delirium refers to confusion, altered awareness, or altered thoughts. It can result from many different illnesses, medications, and other causes.

Delirium was divided into subtypes according to standard DSM-5 criteria: hyperactive delirium, with increased motor activity, loss of control, and restlessness; hypoactive delirium, with decreased activity,

decreased speech, and reduced awareness. Patients with normal psychomotor activity or fluctuating activity levels were classified as having "mixed" delirium.

About 30 percent of patients were diagnosed with delirium on entering palliative care. Of these, the delirium subtype was hyperactive in about 15 percent of patients, hypoactive in 34 percent, and mixed in 51 percent.

Survival time after entering palliative care was shorter for patients with delirium: median 17 days, compared to 28 days for those without delirium. However, the difference was significant only for patients with hypoactive or mixed delirium—with median survival times of 14 and 15 days, respectively.

These differences remained significant after adjustment for other factors. For patients with hyperactive delirium, survival was not different from that in patients without delirium.

While delirium was more common in older patients, the effects on time to death were actually stronger in younger patients. That was consistent with previous studies suggesting shorter survival times in younger patients diagnosed with delirium

Why are different delirium subtypes associated with differing survival times? It may have to do with differences in the underlying causes of and treatment responses. Hyperactive delirium is commonly related to reversible causes, such as medication side effects.

"In contrast, hypoactive delirium is generally related to hypoxia [decreased oxygen levels], metabolic disturbances, and multi-organ failure," Dr. Kim explains. "Therefore, hypoactive delirium could be associated with a higher mortality rate than hyperactive delirium."

Dr. Kim adds, "Also, the earlier mortality in younger patients overturns a conventional assumption for survival prediction of delirium. Although delirium was more prevalent in older patients, as known, the irony is that [delirium](#) predicted shorter survival in younger patients."

Accurate predictions of survival time in [terminally ill patients](#) are important for many reasons—"in terms of ensuring good clinical decision making, developing care strategies, and preparing for the end of life in a dignified manner." The researchers conclude, "Thus, the present findings could facilitate more precise predictions of survival, allowing families to prepare for the patient's death."

**More information:** "Differential Associations Between Delirium and Mortality According to Delirium Subtype and Age: A Prospective Cohort Study." [DOI: 10.1097/PSY.0000000000000239](https://doi.org/10.1097/PSY.0000000000000239)

Provided by Wolters Kluwer Health

Citation: In terminally ill patients, some types of delirium are a sign of 'imminent death' (2015, September 23) retrieved 6 May 2024 from <https://medicalxpress.com/news/2015-09-terminally-ill-patients-delirium-imminent.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---