

# Basketball, soccer, lacrosse lead to most ACL injuries among high school female athletes

October 23 2015

---

A new study finds the overall rate of anterior cruciate ligament (ACL) injuries among high school athletes is significantly higher among females, who are especially likely to experience ACL tears while playing basketball, soccer and lacrosse.

The study, "Sport-Specific Yearly Risk and Incidence of Anterior Cruciate Ligament Tears in High School Athletes: A Systematic Review and Meta-Analysis," will be presented at the 2015 American Academy of Pediatrics Conference & Exhibition in Washington, DC.

Researchers at the Children's Hospital of Philadelphia found significant risk of ACL [injury](#) among both genders, particularly in high-risk sports such as soccer, football, basketball and lacrosse. While the majority of ACL injuries occur in boys, the rate of injury per exposure is higher in girls. Specifically, in girls, the highest ACL injury risks per season were observed in soccer (1.1 percent), basketball (0.9 percent), and lacrosse (0.5 percent). In comparison, the highest risks per season for boys were observed in football (0.8 percent), lacrosse (0.4 percent), and soccer (0.3 percent).

"It has been well established that the risk for ACL tear per athletic exposure is higher in female athletes compared to males," said lead author Alex L. Gornitzky, a fourth-year medical student at the Perelman School of Medicine at the University of Pennsylvania. "As participation rates in high school athletics continues to rise significantly, it has become increasingly important to establish up-to-date, individualized

injury information for high school athletes and their families, who represent a large proportion of patients visiting pediatric orthopaedic and sports medicine clinics."

Knowledge of such sport-specific, seasonal risk is essential for evidenced-based parent-athlete decision-making, accurate physician counseling, and targeted injury-reduction programs for the most at-risk sports, said Mr. Gornitzky, who will present the abstract, "Sport-Specific Yearly Risk and Incidence of Anterior Cruciate Ligament Tears in High School Athletes: A Systematic Review and Meta-Analysis" at 10:53 am on Saturday, Oct. 24 in Saturday, October 24, 2015: 9:50 AM in room 143A of the Walter E. Washington Convention Center.

**More information:** To view the abstract, visit <https://aap.confex.com/aap/2015/webprogrampreliminary/Paper31541.html>.

Provided by American Academy of Pediatrics

Citation: Basketball, soccer, lacrosse lead to most ACL injuries among high school female athletes (2015, October 23) retrieved 18 April 2024 from <https://medicalxpress.com/news/2015-10-basketball-soccer-lacrosse-acl-injuries.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--