

# Cannabis not recommended to prevent post-op nausea

October 14 2015

---



(HealthDay)—Cannabis should not be used to prevent postoperative nausea and vomiting (PONV) because of unacceptable side effects and low effectiveness, according to a study published online Sept. 30 in *Anesthesia & Analgesia*.

Maren Kleine-Brueggeney, M.D., from the University of Bern in Switzerland, and colleagues randomized 40 patients at high risk for PONV to either 0.125 mg/kg intravenous tetrahydrocannabinol (THC) or placebo at the end of surgery before emergence from [anesthesia](#).

The researchers found that the relative risk reduction of overall PONV in the THC group was 12 percent, lower than the clinically significant 25 percent relative risk reduction demonstrated by other drugs used for PONV prophylaxis. After the researchers adjusted for anesthesia time,

the [effect](#) of treatment had an odds ratio of 0.97 (P = 0.97). There were clinically relevant psychotropic THC side effects, mainly consisting of sedation and confusion not related to the effects of anesthesia. The study was discontinued because of THC's inefficacy and the clinically unacceptable side effects.

"Because of an unacceptable side effect profile and uncertain antiemetic effects, IV THC administered at the end of surgery before emergence from anesthesia cannot be recommended for the prevention of PONV in high-risk patients," the authors write.

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Cannabis not recommended to prevent post-op nausea (2015, October 14) retrieved 26 April 2024 from <https://medicalxpress.com/news/2015-10-cannabis-post-op-nausea.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--