


# Study shows chronic fatigue associated with abnormal brain connectivity at rest

October 28 2015

---



Credit: Mary Ann Liebert, Inc., publishers

Patients with chronic fatigue have decreased signaling and communication between specific brain regions when the brain is at rest, and less effective connectivity between these regions strongly correlates with greater fatigue, according to the results of a new study published in *Brain Connectivity*.

Charles Gay, Roland Staud, and colleagues, University of Florida college of Medicine, Gainesville, studied the association between fatigue and altered resting-state connectivity in patients with myalgicencephalomyelitis/chronic fatigue syndrome using functional magnetic resonance imaging.

The researchers used two methods: comparing data on resting-state brain networks; and analyzing [cerebral blood flow](#) in selected brain regions.

**More information:** The article is available free on the [Brain Connectivity](#) website until November 28, 2015.

Provided by Mary Ann Liebert, Inc

Citation: Study shows chronic fatigue associated with abnormal brain connectivity at rest (2015, October 28) retrieved 18 April 2024 from <https://medicalxpress.com/news/2015-10-chronic-fatigue-abnormal-brain-rest.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--