

Drug-resistant *E. coli* continues to climb in community health settings

October 13 2015

Drug-resistant *E. coli* infections are on the rise in community hospitals, where more than half of U.S. patients receive their health care, according to a new study from Duke Medicine.

The study reviewed [patient records](#) at 26 hospitals in the Southeast. By examining [demographic information](#), admission dates and tests, the researchers also found increased antibiotic-resistant infections among [community members](#) who had limited exposure to [health care](#) settings, but who may have acquired the [bugs](#) through some other [environmental factors](#). The findings are published online in *Infection Control & Hospital Epidemiology*.

"We have always considered antibiotic-resistant organisms a problem at large hospitals," said Deverick Anderson, M.D., an infectious disease specialist at Duke University School of Medicine and senior author of the study. "This study goes a long way in demonstrating that the problems with antibiotic-resistant organisms occur in all health care settings, not just large ones. This is also one of the first papers to show these infections are increasing outside of the health care system in the community."

"The larger issue speaks to antibiotics," Anderson said. "Antibiotics are extremely important and useful in medical care, but we know we overuse them."

The study data were gathered through the Duke Infection Control

Network (DICON), which helps [community hospitals](#) and surgery centers across the U.S. prevent infections using education and evidence-based strategies.

The data showed that between 2009 and 2014, the incidence of drug-resistant extended-spectrum beta-lactamase (ESBL)-producing *E. coli* doubled from a rate of 5.28 incidents per 100,000 patients to a rate of 10.5 infections per 100,000. The median age of patients infected with *E. coli* was 72 years.

"Overall, the majority of *E. coli* infections occurred after health care exposure, which makes all hospitals, big and small, important areas of focus to reduce transmission," said lead author Joshua Thaden, M.D., a fellow in Duke's Division of Infectious Diseases. "It's important to consider that a patient's skin may be colonized with drug-resistant bacteria, but because they do not display symptoms, providers may not test them or use extensive contact precautions during care. I think we may be close to a point where it becomes worth the cost and effort to actively screen patients for resistant *E. coli*."

Looking at the timing of patients' infections and when they were last in contact with a health care setting, the researchers also discovered that people with infrequent health care contact were acquiring the superbug at an even faster rate than patients who have regular contact with hospitals or nursing homes. The data showed a greater than three-fold increase in community-associated infections between 2009 and 2014. The sources could be environmental and need further study, Thaden said.

More information: Joshua Thaden, Vance Fowler, Daniel Sexton, Deverick Anderson. "Increasing Incidence of Extended-Spectrum β -Lactamase-Producing *Escherichia coli* in Community Hospitals throughout the Southeastern United States." Web (October 13, 2015).

Provided by Duke University Medical Center

Citation: Drug-resistant E. coli continues to climb in community health settings (2015, October 13) retrieved 5 May 2024 from <https://medicalxpress.com/news/2015-10-drug-resistant-coli-climb-health.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.