

Increased risk of large bowel cancer for each one cm rise in waist circumference

October 27 2015



Credit: UEG

Experts speaking at the 23rd United European Gastroenterology Week (UEG Week 2015) in Barcelona, Spain today revealed compelling

evidence of the link between excess body weight and risk of colorectal cancer (CRC). John Mathers, Professor of Human Nutrition from the Institute of Cellular Medicine at Newcastle University in the UK presented data showing an overall increase of 18% in relative risk of CRC per 5 unit increase in BMI.

"In addition, in men, there is now evidence that increasing waist circumference in middle age is associated with increased bowel cancer risk", says Prof. Mathers. CRC risk was increased by nearly 60% in men who gained at least 10 cm in waist circumference over 10 years. "This increased cancer risk may be due to persistent inflammation in people with obesity".

Patients with Lynch Syndrome (LS) have a higher than normal risk of CRC because of an inherited defect in one of the genes responsible for repairing DNA. Prof. Mathers presented new data showing that, in people with Lynch Syndrome, CRC risk increases with higher [body weight](#) and for those who are obese the risk of CRC is doubled. Quite surprisingly, the increase in CRC risk with higher body weight in people with Lynch Syndrome was about twice as great as that seen in the general population.

Prof. Mathers said "There is now compelling evidence that improved lifestyle, particularly better dietary choices and being more physically active, can help to prevent obesity and this will lower bowel cancer risk". In addition, for those people who are already too heavy, losing weight may reduce their CRC risk but this is an area which requires further study. In his studies with Lynch Syndrome patients, Prof. Mathers observed that aspirin lowered the excess CRC risk seen in patients with obesity, perhaps through its anti-inflammatory effects. "This is a very intriguing finding" said Prof Mathers "which suggests that dietary and other anti-inflammatory agents might be beneficial in reducing CRC risk in people with obesity".

"Bowel cancer is strongly associated with age, obesity and diet - and is driven by inflammation", explains Prof. Mathers. "We can now give the public clear advice on the benefits of staying physically active, eating a healthy diet and avoiding weight gain to lower CRC risk as we get older".

More information: References:

1. Ning Y et al. (2012) *Obesity Reviews* 11, 19-30
2. Song M et al. (2015) *Int. J. Epidemiol.* PMID: 26403814
3. Movahedi M et al. (2015) *J. Clin. Oncol.* PMID: 26282643

Provided by United European Gastroenterology

Citation: Increased risk of large bowel cancer for each one cm rise in waist circumference (2015, October 27) retrieved 6 May 2024 from <https://medicalxpress.com/news/2015-10-large-bowel-cancer-cm-waist.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--