

Nocturnal hypoxemia severity linked to renal RAS activity

October 21 2015



(HealthDay)—In obstructive sleep apnea (OSA), the severity of nocturnal hypoxemia is associated with the extent to which renal renin-angiotensin system (RAS) activity is increased, according to a study published in the Oct. 1 issue of the *American Journal of Respiratory and Critical Care Medicine*.

Ann A. Zalucky, from the University of Calgary in Canada, and colleagues examined the correlation between the severity of nocturnal hypoxemia and RAS activity. In 31 subjects with OSA and 13 obese controls, the authors measured effective renal plasma flow (ERPF) [response](#) to angiotensin II (AngII) challenge, a marker of renal RAS activity.

The researchers found that OSA subjects demonstrated decreased

renovascular sensitivity compared with controls ($P = 0.004$) in response to 60 minutes of AngII challenge. In patients with [severe](#) hypoxemia, the fall in ERPF in response to AngII was less than in those with moderate hypoxemia ($P = 0.001$) and less than in obese control subjects after 30 and 60 minutes of AngII challenge (both P

"The severity of nocturnal hypoxemia influences the magnitude of renal, but not the systemic, RAS activation independently of obesity in patients with OSA," the authors write.

More information: [Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Nocturnal hypoxemia severity linked to renal RAS activity (2015, October 21) retrieved 3 May 2024 from <https://medicalxpress.com/news/2015-10-nocturnal-hypoxemia-severity-linked-renal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
