


Older patients recover more slowly from concussion

October 6 2015



Different working memory activation patterns in healthy control subjects and young and old patients with mild traumatic brain injury. Credit: Radiological Society of North America

Older individuals may have a more difficult time recovering from concussion, according to a new study published online in the journal *Radiology*.

Mild traumatic brain injury (MTBI), or concussion, accounts for 75 percent of all TBI and represents an important public health problem. Difficulty in working memory is frequently reported in [patients](#) after concussion. However, neuropsychological tests, computed tomography (CT) and conventional [magnetic resonance imaging](#) (MRI) generally fail to reveal abnormal findings in these patients. Functional MRI (fMRI) has been increasingly used in assessing patients with MTBI.

"Old age has been recognized as an independent predictor of worse outcome from concussion, but most previous studies were performed on younger adults," said the study's lead author, David Yen-Ting Chen, M.D., from the Department of Radiology at Shuang-Ho Hospital, Taipei Medical University, in New Taipei City, Taiwan.

For this study, Dr. Chen and colleagues used fMRI to evaluate the effect of age on working memory performance and functional activation in the brain after MTBI. The researchers performed fMRI exams on 13 young adults (21-30 years old) and 13 older adults (51-68 years old) with MTBI and 26 age- and gender-matched controls. The first fMRI scan was performed within one month post-injury. A follow-up scan was performed six weeks after the first exam. The researchers then analyzed post-concussion symptoms, neuropsychological test results and working memory activity in both groups.

The analysis revealed that while performing [working memory](#) tasks, the young patients with concussion had initial activation that was greater than normal, known as hyperactivation, compared to young controls in the right precuneus and right inferior parietal gyrus of the brain, whereas the [older patients](#) had hypoactivation (less than normal) compared to older controls in the right precuneus and right [inferior frontal gyrus](#).

In comparing the patients in initial and follow-up study, the young patients had significantly reduced post-concussion symptom score at

follow-up than at the time of the initial exam, but no significant change of the post-concussion symptom score was observed in the older patients, who also showed persistent hypoactivation.

"Taken together, these findings provide evidence for differential neural plasticity across different ages, with potential prognostic and therapeutic implications," said the study's co-author, Ying-Chi Tseng, M.D., from Shuang-Ho Hospital. "The results suggest that MTBI might cause a more profound and lasting effect in older patients."

The researchers hope that these findings might eventually lead to the development of separate management strategies for different age groups following [concussion](#).

More information: "Effect of Age on Working Memory Performance and Cerebral Activation after Mild Traumatic Brain Injury: A Functional MR Imaging Study" *Radiology*, 2015.

Provided by Radiological Society of North America

Citation: Older patients recover more slowly from concussion (2015, October 6) retrieved 11 May 2024 from <https://medicalxpress.com/news/2015-10-older-patients-recover-slowly-concussion.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--