

Opioid overdoses linked to higher prescription rates in British Columbia

October 28 2015



Credit: Flickr, [flickr.com/photos/armydr2008/15910692553/](https://www.flickr.com/photos/armydr2008/15910692553/)

Strong painkillers known as prescription opioids appear to be overprescribed in some regions of British Columbia (B.C.), resulting in higher rates of overdose and death, according to a new study from UBC.


UBC researchers collected data about the amount of opioids prescribed across 79 of the local health [areas](#) of B.C from 2004 to 2013. They found significant variation in prescription rates. Researchers also found that areas with the highest volumes of prescribed opioid purchases were

also the areas that experience the most [overdose deaths](#) involving prescribed [opioid drugs](#).

The study found that, on average, people who live in areas where more [prescription opioids](#) are purchased are not sicker than people who live in areas with low rates of opioid dispensing. The findings point to the need to change prescribing practices and monitor total sales of drugs like fentanyl, oxycodone, morphine, and hydromorphone.

"Our study shows that these [prescription medicines](#) are being used in ways that cause significant harm," said Steve Morgan, a professor in the School of Population and Public Health. "Sometimes the harm is to the person for whom the prescription was written; other times, the harm is to people who appear to have borrowed, bought, or stolen these drugs. We need a solution that addresses both these issues."

Morgan and his colleagues found that many British Columbians (nearly one-third of women and nearly two-thirds of men) who died from a prescribed [opioid overdose](#) did not fill a prescription for such medications in the year prior to their death, indicating that some [prescriptions](#) are being diverted to the illegal drug market.




A map of opioid use in British Columbia, Canada. Credit: Steve Morgan

The researchers calculated that if the higher rates of prescription opioid use in regions such as the southern Okanagan and Fraser Valley were brought down to the lower levels found in North Vancouver, West Vancouver or Richmond, approximately 30 deaths (23 men and 7 women) would have been avoided over the 10-year study period.

Opioid use in B.C. is high compared to the rest of the country but it has been stable for many years and appears better controlled in some areas of the province than others. Morgan believes this is a sign of hope and

that we can save lives.

"With its world-class prescription databases and outstanding medical leaders, B.C. is in an excellent position to monitor prescription practices to avoid fraud and abuse and to improve the management of patients who unfortunately become addicted to opioids," said Morgan. "With adequate investments in monitoring, surveillance, and clinical management, we can ensure that B.C. patients get the treatment they need without the harms."



Rate of opioid-related deaths per 100,000 population.

The study was recently published in *Medical Care*.

More information: *Medical Care*, journals.lww.com/lww-medicalca... iption Opioid.7.aspx

Provided by University of British Columbia

Citation: Opioid overdoses linked to higher prescription rates in British Columbia (2015, October 28) retrieved 27 April 2024 from <https://medicalxpress.com/news/2015-10-opioid-overdoses-linked-higher-prescription.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.