

# Study advances potential test to sort out precancerous pancreatic cysts from harmless ones

November 17 2015

---

In a "look-back" analysis of data stored on 130 patients with pancreatic cysts, scientists at Johns Hopkins have used gene-based tests and a fixed set of clinical criteria to more accurately distinguish precancerous cysts from those less likely to do harm. The findings may eventually help some patients in real time safely avoid surgery to remove harmless cysts. A report on the findings is published in the November issue of *Gastroenterology*.

Currently, doctors must test fluid collected from a needle that penetrates the cyst during a procedure called an endoscopic ultrasound-guided biopsy. The fluid is tested for levels of proteins associated with cancerous cysts, and pathologists look for atypical cells signifying cancer. However, recent studies show that these current methods are accurate only 63 percent of the time, says Anne Marie Lennon, M.D., Ph.D., associate professor of medicine at the Johns Hopkins University School of Medicine and director of the Multidisciplinary Pancreatic Cyst Program.

Building on preliminary gene tests the scientists developed in 2011 that measure the presence of pancreatic cancer-linked genetic markers in cyst fluid samples, the Johns Hopkins-led team expanded the panel of genetic markers and tested them on cyst fluid from 130 patients who had surgery to remove cysts at The Johns Hopkins Hospital and 16 other hospitals in the U.S. and other countries.

In addition to using genetic-based markers, the scientists developed a mathematical model that identifies various types of [pancreatic cysts](#) and their potential for malignancy by taking into account patient characteristics such as age and symptoms, as well as appearance and location of the cysts, based on medical record information from 1,026 patients who had surgery to remove cysts at The Johns Hopkins Hospital.

"The combination of genetic tests and clinical evaluation of patients' cysts may change the way we guide patients on whether surgery to remove cysts is necessary," says Lennon.

To evaluate the validity of the new combination approach, the Johns Hopkins-led team performed genetic tests on the cyst fluid samples and evaluated clinical criteria of the 130 patients.

They were able to identify 12 benign cysts called serous cystadenomas with 98 percent specificity, an indication, they say, of how often a test accurately identifies a disease, and 100 percent sensitivity, a measure of accuracy in correctly identifying people without a disease.

The scientists identified 10 cysts called solid-pseudopapillary neoplasms, which, doctors say, should be surgically removed, with 89 percent sensitivity and 92 percent specificity. Their tests also identified 12 mucinous cystic neoplasms, which doctors often remove, with 97 specificity and 90 percent sensitivity.

Finally, the Johns Hopkins team identified 96 of the more common intraductal papillary mucinous neoplasms with 94 percent sensitivity and 84 percent specificity. The low- and intermediate-risk varieties of these cysts are often monitored carefully when patients have no symptoms. When the malignancy risk level is high, doctors recommend surgical removal.

In the U.S., Lennon says, doctors find pancreatic cysts in approximately 2.6 percent of people who undergo computed tomography scans in the U.S., usually for nonspecific symptoms, such as chronic pain, injury or infection, and three to 13.5 percent of people who undergo magnetic resonance imaging scans. Based on U.S. population statistics for people age 65 and older, this amounts to more than one million people who may face uncertainty about the malignant potential of a pancreatic cyst. Currently, Lennon says approximately 665,600 new pancreatic cysts are identified via CT scan each year in the U.S.

Some harmless cysts should be removed because of their large size or tendency to cause swelling in the organ or jaundice, but Lennon estimates that more than 20 percent of surgically-removed pancreatic cysts thought to be cancerous are later found to be benign.

"Most people who are evaluated for pancreatic cysts are older adults in their 60s and 70s and have comorbidities that can make surgery more risky," says Lennon. "We want to avoid unnecessary surgery whenever we can." The complications include fistulas, infections, and some deaths, which also occur.

The researchers caution that, 40 percent of the time, doctors are unable to collect enough cyst fluid for standard analysis, a problem they say their genetic testing method solves by requiring only very small samples of cystic DNA. But they caution that cyst fluid removal is not risk-free and can cause bleeding, infection and inflammation in a very small percentage of patients.

To determine whether DNA changes present in the cyst could be accurately detected before surgery, Lennon and her colleagues compared genetic tests on cyst fluid samples obtained during endoscopic ultrasound before surgery and from cyst specimens surgically removed in 24 of the 130 study subjects. They found a similar number of genetic changes in

21 of the samples before surgery and 20 of the surgically removed cyst specimens.

"Our study shows that our expanded panel of genetic markers, combined with clinical acumen, can accurately differentiate between types of pancreatic cysts and identify cysts that are safe to watch over time and cysts that need to be removed," says Ralph Hruban, M.D., Baxley Professor in Pathology and director of the pathology department for the Johns Hopkins University School of Medicine and director of the Sol Goldman Pancreatic Cancer Research Center.

"Correctly identifying whether pancreatic cysts pose a high risk for malignancy is key to preventing an often deadly cancer, or avoiding a needless surgery for cysts not likely to be a problem," says Bert Vogelstein, M.D., Clayton Professor of Oncology and director of the Ludwig Center at the Johns Hopkins Kimmel Cancer Center. He was also a co-author of the study.

Lennon says the team plans to study the combination approach on more varieties of pancreatic cysts, especially the rarer types. To further validate the findings, they must test the approach on larger numbers of patients, determine how well the approach identifies cysts that need to be removed in newly-diagnosed patients, and evaluate its cost effectiveness.

**More information:** Simeon Springer et al. A Combination of Molecular Markers and Clinical Features Improve the Classification of Pancreatic Cysts, *Gastroenterology* (2015). [DOI: 10.1053/j.gastro.2015.07.041](https://doi.org/10.1053/j.gastro.2015.07.041)

Provided by Johns Hopkins University School of Medicine

Citation: Study advances potential test to sort out precancerous pancreatic cysts from harmless ones (2015, November 17) retrieved 9 May 2024 from

<https://medicalxpress.com/news/2015-11-advances-potential-precancerous-pancreatic-cysts.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.