

Researchers present findings on cardiac risks for patients with chronic kidney disease

November 8 2015

Chronic Kidney Disease (CKD), which afflicts more than 26 million Americans, is a condition in which individuals experience a slow loss of kidney function over time. At the American Heart Association Scientific Sessions 2015, researchers from the Perelman School of Medicine at the University of Pennsylvania will present findings from their analysis of the Chronic Renal Insufficiently Cohort (CRIC) study to evaluate risk markers for adverse cardiac events in patients with CKD.

Researchers examined data from the CRIC study to determine how the epidemiology of [peripheral artery disease](#) (PAD) in patients with CKD differs according to patient's age and gender. Data revealed that the [risk](#) of PAD is higher in women as compared to men. Results also showed that while age affected the risk of PAD in men more directly, the risk in women was consistent from year to year, and it began earlier in life. But by the time both men and women reached their 70s, the rate of risk was relatively similar.

"We set out to determine whether there were gender differences in peripheral artery disease in patients with CKD, and found that women had an increased risk compared to men; however, this relationship was modified by age," said the study's lead author, Grace Wang, MD, an assistant professor of Surgery and Radiology, and director of the Vascular Laboratory at the Hospital of the University of Pennsylvania. "The analysis showed that women were more at risk for PAD early on, starting around age 40, but the risk in men became greater as they aged."

Results showed that the risk of PAD in women was 1.5 times higher than the risk in men, prior to age 70. Overall, women with CKD have a higher PAD incidence at nearly 23 percent, as compared to an almost 14 percent in men. Researchers also noted that additional studies should be conducted to understand the impact of earlier detection of PAD in [women](#), as a means for preventing or proactively treating peripheral artery disease.

Provided by University of Pennsylvania School of Medicine

Citation: Researchers present findings on cardiac risks for patients with chronic kidney disease (2015, November 8) retrieved 13 March 2024 from <https://medicalxpress.com/news/2015-11-cardiac-patients-chronic-kidney-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--