

# Dental implants frequently lead to complications

#### November 5 2015

Almost 8 percent of patients experience loss of at least one implant within ten years. Even more develop peri-implantitis. Patients with periodontitis run a greater risk of both implant loss and peri-implantitis. A doctoral thesis at Sahlgrenska Academy has explored the various issues.

Approximately 30,000 Swedes receive dental implant therapy every year. The method is common in connection with partial or full edentulism (toothlessness), and Sweden is among the countries with the most <u>dental implants</u> per capita.

# **Biological complications**

Previous evaluations have frequently been limited to <u>observational</u> <u>studies</u> of patient populations in specialist dental care. Few attempts have been made to analyze biological complications such as peri-implantitis, which is characterized by inflammation in adjacent tissues and loss of supporting jawbone.

A national project evaluated the therapy in terms of patient-reported outcomes, implant loss and prevalence of peri-implantitis.

#### High degree of satisfaction

"The results of the initial questionnaire survey, which included 4,716



randomly selected <u>patients</u>, indicated a high degree of satisfaction, particularly among men and older respondents," says Jan Derks, researcher at Sahlgrenska Academy.

### **Developed peri-implantitis**

A total of 596 patients at 37 Swedish clinics were examined at a 9-year follow-up.

"Altogether, 7.6 % of patients had lost at least one implant and 14.5 % had developed peri-implantitis with pronounced bone loss," Dr. Derks says.

Patients suffering from periodontitis exhibited an increased risk of severe peri-implantitis, while both periodontitis and smoking were associated with early implant loss.

# **Progresses quickly**

"Peri-implantitis appears to develop within a few years and then progresses quickly and at an accelerating pace," Dr. Derks says.

The results of the study provide crucial input to both patients and dental practitioners in the effort to minimize the risk of implant loss and periimplantitis.

More information: Full thesis: <a href="mailto:gupea.ub.gu.se/handle/2077/39544">gupea.ub.gu.se/handle/2077/39544</a>

Provided by University of Gothenburg



Citation: Dental implants frequently lead to complications (2015, November 5) retrieved 3 April 2024 from <a href="https://medicalxpress.com/news/2015-11-dental-implants-frequently-complications.html">https://medicalxpress.com/news/2015-11-dental-implants-frequently-complications.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.