

Study examines factors that may contribute to hypersexuality

November 4 2015

In a study of European men, hypersexuality—a preoccupation with sexual fantasy or an excessive indulgence in sexual activity—correlated with proneness to sexual boredom and problems with erectile function.

The study, which included 911 Croatian and 210 German men who were currently in a relationship, contributes to existing knowledge on hypersexuality. The findings indicate that therapies for hypersexual men should include sex therapy principals that may enhance erectile functioning and address sexual boredom.

"In some [men](#), hypersexual behavior may serve as a coping mechanism for sexual boredom. In addition, [erectile dysfunction](#) within an intimate relationship can accompany this simultaneously," said Verena Klein, lead author of *The Journal of Sexual Medicine* study.

More information: Verena Klein et al. Erectile Dysfunction, Boredom, and Hypersexuality among Coupled Men from Two European Countries, *The Journal of Sexual Medicine* (2015). [DOI: 10.1111/jsm.13019](#)

Provided by Wiley

Citation: Study examines factors that may contribute to hypersexuality (2015, November 4) retrieved 24 April 2024 from

<https://medicalxpress.com/news/2015-11-factors-contribute-hypersexuality.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.