

Mixed martial arts bloodier but less dangerous than boxing, study finds

November 5 2015



Mixed martial arts fighters Cody Krahn (left) and Curtis Demarce spar while fellow fighter Victor Valimaki looks on. Credit: University of Alberta

Mixed martial arts has a reputation for being one of the most brutal and bloody of all contact sports, but the reality is boxing poses a greater risk of serious injury, according to new research from the University of Alberta.

Researchers at the U of A's Glen Sather Sports Medicine Clinic

reviewed a decade's worth of data from medical examinations following mixed martial arts and boxing matches and found that MMA fighters face a slightly higher risk of minor injuries. Boxers, however, are more likely to experience serious harm from concussions and other head trauma, loss of consciousness, [eye injuries](#), smashed noses and broken bones.

"Yes, you're more likely to get injured if you're participating in mixed martial arts, but the injury severity is less overall than boxing," explained lead author Shelby Karpman, a [sports medicine](#) physician at the Glen Sather clinic. "Most of the blood you see in mixed martial arts is from bloody noses or facial cuts; it doesn't tend to be as severe but looks a lot worse than it actually is."

Research from ringside

The research offers a first-of-its-kind glimpse into the dangers of the two combative sports in Canada, and is the direct result of Karpman's quarter-century of experience as a ringside physician conducting post-fight exams, which are mandatory in both sports.

In this study, Karpman and U of A Faculty of Rehabilitation Medicine colleagues Leah Phillips, Ziling Qin and Doug Gross, and Patrick Reid of the Edmonton Combative Sports Commission, reviewed post-fight records from 1,181 MMA fighters and 550 boxers who fought matches in Edmonton between 2003 and 2013.

They found that 59.4 per cent of MMA fighters suffered some form of injury in their bouts—significantly higher than the injury rate of 49.8 per cent for boxers. Most of these injuries were bruises and contusions. But boxers were more likely to experience loss of consciousness during the bout (7.1 per cent compared with 4.2 per cent for MMA fighters) or serious eye injuries.

Boxers were also significantly more likely to receive medical suspensions due to injuries suffered during bouts.

Karpman said there is risk in any contact sport but that MMA, more than any other, faces a stigma from the medical community from physicians who see the sport as ultra-bloody and violent. As a result, fighters have become "an undertreated athletic population," and these research results should help them understand the risks of climbing into the ring, he said.

"These guys do not get the respect they deserve for what they're doing—or the medical treatment—because the [medical community](#) doesn't want to deal with such a bloody sport with head injuries and concussions," Karpman said.

Most fighters understand the risks they face before they get into the ring, but the research results do line up with the real-life experience of 14-year MMA veteran Victor Valimaki.

"There are definitely risks. I've been pretty messed up," said Valimaki, who says he's suffered just about every injury imaginable: broken bones—both feet and arms, an ankle, collar bone—a busted orbital bone, and numerous knee and shoulder joint issues.

"Most injuries happen during training. Injuries during an actual fight are superficial—typically black eyes, cuts and the odd broken hand," he said.

Banning combative sports not the answer

Karpman says it's puzzling that in a sport like hockey, catastrophic blows to the head can land former NHL players like Scott Stevens and Chris Pronger in the hall of fame, whereas MMA and boxing are vilified with frequent calls to ban the sport with little understanding of the true risks.

"I always say if you're going to ban a sport, you need statistics. Just watching mixed [martial arts](#) twice on TV does not cut it. And even if you ban a sport, you're not going to stop it. You're just going to take it underground where they're not going to receive medical care."

The research was published in the peer-reviewed *Clinical Journal of Sports Medicine*.

Provided by University of Alberta

Citation: Mixed martial arts bloodier but less dangerous than boxing, study finds (2015, November 5) retrieved 5 May 2024 from <https://medicalxpress.com/news/2015-11-martial-arts-bloodier-dangerous.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--